



THE ORDER OF APPEARANCE OF THE PRODUCT OF FIVE CONSECUTIVE LUCAS NUMBERS

DIEGO MARQUES — PAVEL TROJOVSKÝ

ABSTRACT. Let F_n be the n th Fibonacci number and let L_n be the n th Lucas number. The order of appearance $z(n)$ of a natural number n is defined as the smallest natural number k such that n divides F_k . For instance, $z(F_n) = n = z(L_n)/2$ for all $n > 2$. In this paper, among other things, we prove that

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) = \frac{n(n+1)(n+2)(n+3)(n+4)}{12}$$

for all positive integers $n \equiv 0, 8 \pmod{12}$.

1. Introduction

Let $(F_n)_n$ be the Fibonacci sequence given by $F_{n+2} = F_{n+1} + F_n$, for $n \geq 0$, where $F_0 = 0$ and $F_1 = 1$. Let $(L_n)_n$ be the Lucas sequence which follows the same recursive pattern as the Fibonacci numbers, but with initial values $L_0 = 2$ and $L_1 = 1$. These numbers are well-known for possessing amazing properties (for example consult [4]). The period $k(m)$ of the Fibonacci sequence modulo a positive integer m is the smallest positive integer n such that

$$F_n \equiv 0 \pmod{m} \quad \text{and} \quad F_{n+1} \equiv 1 \pmod{m}.$$

The study of the divisibility properties of Fibonacci numbers has always been a popular area of research. Let n be a positive integer, the *order (or rank) of appearance* of n in the Fibonacci sequence, denoted by $z(n)$, is defined as the smallest positive integer k , such that $n \mid F_k$ (some authors also call it *order of apparition*, or *Fibonacci entry point*). There are several results about $z(n)$ in the literature. For example, $z(m) \leq 2m$, for all $m \geq 1$ (see [15] and [10] for improvements) and in the case of a prime number p , one has the better upper bound $z(p) \leq p + 1$, which is a consequence of the known congruence

© 2014 Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: 11B39.

Keywords: Fibonacci numbers, Lucas numbers, order of appearance, p -adic order.

Research supported in part by DPP-UnB, FAP-DF, CNPq and IPCZ.

$F_{p-\left(\frac{p}{5}\right)} \equiv 0 \pmod{p}$ for $p \neq 2, 5$, where $\left(\frac{a}{q}\right)$ denotes the Legendre symbol of a with respect to a prime $q > 2$. We will use Pochhammer polynomial $n^{(k)} = n(n+1)(n+2)\dots(n+k-1)$ for the simplification of notation in the following text.

In recent papers, the first author [5]–[9] found explicit formulas for the order of appearance of integers related to Fibonacci and Lucas numbers, such as $C_m \pm 1, C_n C_{n+1} C_{n+2} C_{n+3}$ and C_n^k , where C_n represents F_n or L_n .

In this paper, we continue this program by studying the order of appearance of the product of five consecutive Lucas numbers. Our main result is the following.

THEOREM 1.1. *Let n be any nonnegative integer. Then*

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) = \begin{cases} n^{(5)}, & n \equiv 1 \pmod{6}; \\ \frac{1}{2}n^{(5)}, & n \equiv 2, 10, 14, 18, 22, 30, 34 \pmod{36}; \\ \frac{1}{3}n^{(5)}, & n \equiv 3, 5 \pmod{6}; \\ \frac{1}{4}n^{(5)}, & n \equiv 4 \pmod{12}; \\ \frac{1}{6}n^{(5)}, & n \equiv 6, 26 \pmod{36}; \\ \frac{1}{12}n^{(5)}, & n \equiv 0, 8 \pmod{12}. \end{cases} \tag{1.1}$$

Remark 1. The completeness of cases in Theorem 1.1 follows from the fact that the first case and the third case together include all positive odd integers n and the other cases include all nonnegative even integers n .

2. Auxiliary results

Before proceeding further, we recall some facts on Fibonacci numbers for the convenience of the reader.

The p -adic valuation (or order) of r , $\nu_p(r)$, is the exponent of the highest power of a prime p which divides r . The p -adic order of the Fibonacci and Lucas numbers was completely characterized, see [14], [16] and [12]. For instance, from the main results of Lengyel [12], we extract the following result.

LEMMA 2.1. *Let p be any prime. For $n \geq 1$, we have*

$$\nu_2(F_n) = \begin{cases} 0 & \text{if } n \equiv 1, 2 \pmod{3}; \\ 1 & \text{if } n \equiv 3 \pmod{6}; \\ 3 & \text{if } n \equiv 6 \pmod{12}; \\ \nu_2(n) + 2 & \text{if } n \equiv 0 \pmod{12}, \end{cases}$$

$$\nu_5(F_n) = \nu_5(n),$$

and for any prime $p \neq 5$ and $p > 2$

$$\nu_p(F_n) = \begin{cases} \nu_p(n) + \nu_p(F_{z(p)}) & \text{if } n \equiv 0 \pmod{z(p)}; \\ 0 & \text{if } n \not\equiv 0 \pmod{z(p)}. \end{cases}$$

LEMMA 2.2. *Let p be any prime, let $k(p)$ be the period modulo p of the Fibonacci sequence. For $n \geq 1$, we have*

$$\nu_2(L_n) = \begin{cases} 0, & n \equiv 1, 2 \pmod{3}; \\ 2, & n \equiv 3 \pmod{6}; \\ 1, & n \equiv 0 \pmod{6}, \end{cases}$$

and for any prime $p > 2$,

$$\nu_p(L_n) = \begin{cases} \nu_p(n) + \nu_p(F_{z(p)}), & k(p) \neq 4z(p) \text{ and } n \equiv \frac{z(p)}{2} \pmod{z(p)}; \\ 0, & \text{otherwise.} \end{cases}$$

Remark 2. Since $k(5) = 20$ and $z(5) = 5$, we have $k(5) = 4z(5)$ and so the previous lemma yields $\nu_5(L_n) = 0$. In fact, the same happens for the primes

13, 17, 37, 53, 61, 73, 89, 97, 109, 113, 137, 149, 157, 173, 193, 197, ...

which is the OEIS sequence A053028. We point out an interesting result of Lagarias [11] concerning the density of this set of primes.

LEMMA 2.3 (Cf. Lemma 2.1 [9]). *We have*

- (a) $F_n \mid F_m$ if and only if $n \mid m$.
- (b) $L_n \mid F_m$ if and only if $n \mid m$ and m/n is even.
- (c) $L_n \mid L_m$ if and only if $n \mid m$ and m/n is odd.
- (d) $F_{2n} = F_n L_n$.
- (e) $\gcd(L_n, L_{n+1}) = \gcd(L_n, L_{n+2}) = 1$.

LEMMA 2.4 (Cf. Lemma 2.2 of [9]). *We have*

- (a) If $F_n \mid m$, then $n \mid z(m)$.
- (b) If $L_n \mid m$, then $2n \mid z(m)$.
- (c) If $n \mid F_m$, then $z(n) \mid m$.

LEMMA 2.5. *Let k, n, m be any positive integers. We have*

- (a) If $n \equiv 0, 3 \pmod{6}$, then $2F_n \mid F_{2n}$.
If $n \equiv 2 \pmod{4}$, then $3F_n \mid F_{2n}$.
- (b) If $n \equiv 1, 2 \pmod{3}$, then $2F_n \mid F_{3n}$.
- (c) If $n \equiv 6 \pmod{12}$, then $6F_n \mid F_{2n}$.
- (d) If $n \equiv 2 \pmod{4}$, then $6F_n \mid F_{6n}$.

- (e) If $m \nmid F_{kn}$, then $m \nmid F_n$.
- (f) If $n \equiv 2, 10, 14, 18, 22, 30 \pmod{36}$, then $L_n L_{n+4} \nmid F_{\frac{1}{3}n^{(5)}}$.
- (g) If $n \equiv 2 \pmod{4}$, then $L_{n+2} \nmid F_{\frac{1}{4}n^{(5)}}$.
- (h) If $1 \leq k \leq 5$, then $\nu_2\left(\prod_{i=0}^k L_{n+i}\right) \leq 3$.

Proof.

- (a) Using the identity $F_{2n} = F_n L_n$ and Lemma 2.2, we clearly obtain the assertion.
- (b) Using the identity $F_{3n} = F_n(L_{2n} + (-1)^n)$, see [4, p. 92], the fact that L_{2n} is odd for $n \equiv 1, 2 \pmod{3}$ by Lemma 2.2 we clearly obtain the assertion.
- (c) Using the identity $F_{2n} = F_n L_n$ and Lemma 2.2 we obtain the assertion.
- (d) Using the identity $F_{6n} = F_{3n} L_{3n} = L_{3n} F_n(L_{2n} + (-1)^n)$, see [4, p. 92], and the fact that $6 \mid L_{3n}$ for $n \equiv 2 \pmod{4}$, with respect to Lemma 2.2, we have the assertion.
- (e) Let us consider that $m \mid F_n$. Using the well-known property $F_n \mid F_{kn}$ we obtain $m \mid F_{kn}$.
- (f) To prove the assertion it is suffice to show that $\nu_3(L_n L_{n+4}) > \nu_3(F_{\frac{1}{3}n^{(5)}})$ for $n \equiv 2, 10, 14, 18, 22, 30 \pmod{36}$ (thus $n \equiv 2 \pmod{4}$ and $n \not\equiv 6, 26, 34 \pmod{36}$). Using Lemmas 2.1, 2.2 and the clear fact that $4 \mid \frac{1}{3}n^{(5)}$ for any nonnegative integer n we obtain

$$\nu_3(L_n) = \begin{cases} \nu_3(n) + 1, & n \equiv 2 \pmod{4}; \\ 0, & n \not\equiv 2 \pmod{4}, \end{cases} \quad (2.1)$$

$$\nu_3(F_n) = \begin{cases} \nu_3(n) + 1, & n \equiv 0 \pmod{4}; \\ 0, & n \not\equiv 0 \pmod{4}, \end{cases}$$

hence

$$\begin{aligned} \nu_3(L_n L_{n+4}) &= \nu_3(L_n) + \nu_3(L_{n+4}) \\ &= (\nu_3(n) + 1) + (\nu_3(n+4) + 1) \\ &= \nu_3(n) + \nu_3(n+4) + 2 \end{aligned}$$

and

$$\begin{aligned} \nu_3(F_{\frac{1}{3}n^{(5)}}) &= \nu_3\left(\frac{1}{3}n^{(5)}\right) + 1 = \nu_3(n^{(5)}) \\ &= \nu_3(n) + \nu_3(n+4) + 1, \end{aligned}$$

as clearly $\nu_3((n+1)(n+2)(n+3)) = 1$ holds for $n \not\equiv 6, 7, 8 \pmod{9}$ and all cases from the assertion are in this form.

(g) For $n \equiv 2 \pmod{4}$ we have $\frac{1}{4}n^{(5)}/(n+2) \equiv 1 \pmod{2}$, hence the assertion follows from Lemma 2.3 (b).

(h) Since there are unique ϵ and δ belonging to $\{0, \dots, 5\}$ such that

$$n + \epsilon \equiv 3 \pmod{6} \quad \text{and} \quad n + \delta \equiv 0 \pmod{6},$$

we have

$$\nu_2 \left(\prod_{i=0}^k L_{n+i} \right) \leq \sum_{i=0}^5 \nu_2(L_{n+i}) = \nu_2(L_{n+\epsilon}) + \nu_2(L_{n+\delta}) = 2 + 1 = 3.$$

Thus the lemma follows. □

Remark 3. The reader may be wondering why this paper deals with Lucas numbers, but it does not study the Fibonacci case. The reason is exactly that the previous item (h) does not hold for Fibonacci numbers. Actually, $\nu_2 \left(\prod_{i=0}^k F_{n+i} \right)$ can be sufficiently large which causes the substantial increasing in the number of cases to be studied.

3. The proof of Theorem 1.1

Since there are at least two even numbers among $n, n+1, n+2, n+3, n+4$, we conclude (using Lemma 2.3 (b)) that

$$L_{n+i} \mid F_{n^{(5)}} \quad \text{for } i = 0, 1, 2, 3, 4. \tag{3.1}$$

We will consider these cases:

- Let $n \equiv 1 \pmod{6}$. Then $\gcd(L_n, L_{n+3}) = \gcd(L_n, L_{n+4}) = 1$. This together with Lemma 2.3 (e) implies that the numbers $L_n, L_{n+1}, L_{n+2}, L_{n+3}$ and L_{n+4} are pairwise coprime. Thus (3.1), together with Lemma 2.4 (c), leads to

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid n^{(5)}. \tag{3.2}$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$ with respect to Lemma 2.4 (b). Since $n, \frac{n+1}{2}, 2(n+2), \frac{n+3}{2}, n+4$ are pairwise coprime, then

$$2n \frac{n+1}{2} 2(n+2) \frac{n+3}{2} (n+4) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \tag{3.3}$$

Combining (3.2) and (3.3)

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) = n^{(5)}.$$

- Let $n \equiv 4 \pmod{12}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{4}n^{(5)}} \quad (3.4)$$

for $i = 0, 1, 2, 3, 4$. Further $\gcd(L_n, L_{n+3}) = \gcd(L_n, L_{n+4}) = 1$. This together with Lemma 2.3 (e) yields that the numbers $L_n, L_{n+1}, L_{n+2}, L_{n+3}$, and L_{n+4} are pairwise coprime. Thus Lemma 2.4 (c) implies that

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid \frac{1}{4}n^{(5)}. \quad (3.5)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Since $n/2, n+1, (n+2)/2, n+3, (n+4)/2$ are pairwise coprime, then

$$2\frac{n}{4}(n+1)\frac{n+2}{2}(n+3)\frac{n+4}{2} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \quad (3.6)$$

Thus, combining (3.5) and (3.6) we have

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{8}n^{(5)}, \frac{1}{4}n^{(5)} \right\}.$$

Now, we show that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \nmid F_{\frac{1}{8}n^{(5)}}.$$

In fact, by using Lemma 2.3 (b) we have

$$\begin{aligned} L_n &\nmid F_{\frac{1}{8}n^{(5)}} \quad \text{for } n \equiv 16 \pmod{24}, \\ L_{n+4} &\nmid F_{\frac{1}{8}n^{(5)}} \quad \text{for } n \equiv 4 \pmod{24}. \end{aligned}$$

- Let $n \equiv 8 \pmod{12}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{12}n^{(5)}} \quad (3.7)$$

for $i = 0, 1, 2, 3, 4$. Further $\gcd(L_n, L_{n+3}) = \gcd(L_n, L_{n+4}) = 1$ and together with Lemma 2.3 (e), we observe that the numbers $L_n, L_{n+1}, L_{n+2}, L_{n+3}$ and L_{n+4} are pairwise coprime. Thus Lemma 2.4 (c) implies that

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid \frac{1}{12}n^{(5)}. \quad (3.8)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Observe that there exist $a, b, c, d \in \{0, 1\}$, with $a+b = c+d = 1$, such that $n/4^a, (n+1)/3^c, (n+2)/2, n+3, (n+4)/(4^b \cdot 3^d)$ are pairwise coprime, then

$$\frac{1}{24}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2. \quad (3.9)$$

Thus, using (3.8) and (3.9) we have

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) = \frac{1}{12}n^{(5)}.$$

- Let $n \equiv 0 \pmod{12}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{12}n^{(5)}} \quad (3.10)$$

for $i = 0, 1, 2, 3, 4$. Further $\gcd\left(\frac{L_n}{2}, L_{n+3}\right) = 1$ and $\gcd(L_n, L_{n+4}) = 1$. Hence using Lemma 2.5 (a) we obtain

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 2F_{\frac{1}{12}n^{(5)}} \mid F_{\frac{1}{6}n^{(5)}}$$

and

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid \frac{1}{6}n^{(5)}. \quad (3.11)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Observe that there exist $a, b, c, d \in \{0, 1\}$, with $a+b = c+d = 1$, such that $n/(4^a \cdot 3^c)$, $n+1$, $(n+2)/2$, $(n+3)/3^d$, $(n+4)/4^b$ are pairwise coprime, then

$$\frac{1}{24}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2 \quad (3.12)$$

and therefore

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{12}n^{(5)}, \frac{1}{6}n^{(5)} \right\}.$$

We show that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{12}n^{(5)}}.$$

The proof will be based on comparing p -adic orders of

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \quad \text{and} \quad F_{\frac{1}{12}n^{(5)}} \quad \text{for all primes } p.$$

Thus we shall prove that

$$\nu_p(F_{\frac{1}{12}n^{(5)}}) \geq \nu_p(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \quad (3.13)$$

holds for all primes p .

Using Lemma 2.5 (h), Lemma 2.1 and the clear fact that

$$\frac{1}{12}n^{(5)} \equiv 0 \pmod{48},$$

we have

$$\begin{aligned} \nu_2(F_{\frac{1}{12}n^{(5)}}) &= \nu_2\left(\frac{1}{12}n^{(5)}\right) + 2 \\ &= \nu_2(n^{(5)}) - 2 + 2 \geq 6 \\ &> \nu_2(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \end{aligned}$$

Now, we will consider $p \neq 2$. Suppose that $\nu_p(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \neq 0$ (otherwise the desired inequality is directly proved). Since $\frac{L_n}{2}$, L_{n+1} , L_{n+2} , L_{n+3} and L_{n+4} are pairwise coprime, p divides only one of L_n , L_{n+1} , L_{n+2} , L_{n+3} , L_{n+4} , say $p \mid L_{n+\delta}$ for some $\delta \in \{0, 1, 2, 3, 4\}$. Thus $p \mid L_{n+\delta} \mid F_{\frac{1}{12}n^{(5)}}$

implying $z(p) \mid \frac{1}{12}n^{(5)}$, by Lemma 2.4 (c). Therefore using Lemma 2.1 and Lemma 2.2 we obtain

$$\begin{aligned} \nu_p(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) &= \nu_p(L_{n+\delta}) \\ &\leq \nu_p(n + \delta) + \nu_p(F_{z(p)}) \\ &\leq \nu_p\left(\frac{1}{12}n^{(5)}\right) + \nu_p(F_{z(p)}) \\ &= \nu_p\left(F_{\frac{1}{12}n^{(5)}}\right). \end{aligned}$$

- Let $n \equiv 3 \pmod{6}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{3}n^{(5)}} \quad \text{for } i = 0, 1, 2, 3, 4. \quad (3.14)$$

Further $\gcd(\frac{L_n}{2}, L_{n+3}) = \gcd(L_n, L_{n+4}) = \gcd(L_{n+1}, L_{n+4}) = 1$ and together with Lemma 2.3 (e) the numbers $\frac{L_n}{2}, L_{n+1}, L_{n+2}, L_{n+3}$ and L_{n+4} are pairwise coprime. Hence using Lemma 2.5 (a) we obtain

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 2F_{\frac{1}{3}n^{(5)}} \mid F_{2\frac{1}{3}n^{(5)}}.$$

In particular,

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid \frac{2}{3}n^{(5)}. \quad (3.15)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Observe that there exist $a, b, c, d \in \{0, 1\}$, with $a + b = c + d = 1$, such that $n/3^a, (n+1)/2^c, n+2, (n+3)/(3^b \cdot 2^d), n+4$ are pairwise coprime, then

$$\frac{1}{6}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2. \quad (3.16)$$

Thus, using (3.15) and (3.16) we have

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{3}n^{(5)}, \frac{2}{3}n^{(5)} \right\}.$$

We show that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{3}n^{(5)}}.$$

The proof will be again based on comparing p -adic orders of

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \quad \text{and} \quad F_{\frac{1}{3}n^{(5)}}$$

for all primes p . Thus we prove that

$$\nu_p\left(F_{\frac{1}{3}n^{(5)}}\right) \geq \nu_p(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \quad (3.17)$$

holds for all primes p . This relation clearly holds for $p = 5$ with respect to Lemma 2.1 and Lemma 2.2. Using Lemma 2.5 (h), Lemma 2.1 and

the clear fact that $\frac{1}{3}n^{(5)} \equiv 0 \pmod{24}$ we obtain

$$\begin{aligned} \nu_2(F_{\frac{1}{3}n^{(5)}}) &= \nu_2\left(\frac{1}{3}n^{(5)}\right) + 2 = \nu_2(n^{(5)}) + 2 \\ &\geq 5 > \nu_2(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \end{aligned}$$

When $p \neq 2$ and $p \neq 5$, the proof of (3.17) can be done by the same way as in the case $n \equiv 0 \pmod{12}$.

- Let $n \equiv 5 \pmod{6}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{3}n^{(5)}} \quad \text{for } i = 0, 1, 2, 3, 4.$$

Further $\gcd(L_n, L_{n+3}) = \gcd(L_n, L_{n+4}) = \gcd(\frac{L_{n+1}}{2}, L_{n+4}) = 1$, and together with Lemma 2.3 (e) the numbers $L_n, \frac{L_{n+1}}{2}, L_{n+2}, L_{n+3}$ and L_{n+4} are pairwise coprime. Hence using Lemma 2.5 (b) we obtain

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 2F_{\frac{1}{3}n^{(5)}} \mid F_{3\frac{1}{3}n^{(5)}}$$

and then

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid n^{(5)}. \quad (3.18)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Observe that there exist $a, b, c, d \in \{0, 1\}$ with $a+b=c+d=1$, such that $n, (n+1)/(2^a \cdot 3^b), n+2, (n+3)/2^b, (n+4)/3^b$ are pairwise coprime, then

$$\frac{1}{6}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2. \quad (3.19)$$

Thus, using (3.18) and (3.19) we have

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{3}n^{(5)}, \frac{2}{3}n^{(5)}, n^{(5)} \right\}.$$

The fact that $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{3}n^{(5)}}$ holds can be proved in the same way as in the case $n \equiv 3 \pmod{6}$.

- Let $n \equiv 6 \pmod{36}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{6}n^{(5)}} \quad \text{for } i = 0, 1, 2, 3, 4.$$

Further $\gcd(\frac{L_n}{2}, L_{n+3}) = \gcd(\frac{L_n}{2}, \frac{L_{n+4}}{3}) = \gcd(L_{n+1}, \frac{L_{n+4}}{3}) = 1$ and together with Lemma 2.3 (e) the numbers $\frac{L_n}{2}, L_{n+1}, L_{n+2}, L_{n+3}$ and $\frac{L_{n+4}}{3}$ are pairwise coprime. Hence using Lemma 2.5 (c) we obtain

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 6F_{\frac{1}{6}n^{(5)}} \mid F_{\frac{1}{3}n^{(5)}}$$

and

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid \frac{1}{3}n^{(5)}. \quad (3.20)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Since $n/6, n+1, n+2, n+3, (n+4)/2$ are pairwise coprime, we have that

$$\frac{1}{12}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2. \quad (3.21)$$

Thus, using (3.20) and (3.21) yields

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{6}n^{(5)}, \frac{1}{3}n^{(5)} \right\}.$$

So, it remains to prove that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{6}n^{(5)}}.$$

Using Lemma 2.5 (h) and the clear fact $\frac{1}{6}n^{(5)} \equiv 0 \pmod{12}$ by Lemma 2.1, we get

$$\begin{aligned} \nu_2 \left(F_{\frac{1}{6}n^{(5)}} \right) &= \nu_2 \left(\frac{1}{6}n^{(5)} \right) + 2 = \nu_2(n^{(5)}) + 1 \\ &\geq 5 > \nu_2(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \end{aligned}$$

For $p > 2$ we can prove that $\nu_p(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \leq \nu_p(F_{\frac{1}{6}n^{(5)}})$ in the same way as in the case $n \equiv 0 \pmod{12}$.

- Let $n \equiv 26 \pmod{36}$. Using Lemma 2.3 (b) we clearly have that

$$L_{n+i} \mid F_{\frac{1}{6}n^{(5)}} \quad \text{for } i = 0, 1, 2, 3, 4.$$

Further $\gcd(\frac{L_n}{3}, L_{n+3}) = \gcd(\frac{L_n}{3}, \frac{L_{n+4}}{2}) = \gcd(L_{n+1}, \frac{L_{n+4}}{2}) = 1$ and this together with Lemma 2.3 (e) implies that the numbers $\frac{L_n}{3}, L_{n+1}, L_{n+2}, L_{n+3}$ and $\frac{L_{n+4}}{2}$ are pairwise coprime. Hence using Lemma 2.5 (d) we obtain

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 6F_{\frac{1}{6}n^{(5)}} \mid F_{n^{(5)}}$$

and

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid n^{(5)}. \quad (3.22)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Since $n/2, n+1, n+2, n+3, (n+4)/6$ are pairwise coprime, we have that

$$\frac{1}{12}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})/2. \quad (3.23)$$

Thus, (3.22) and (3.23) yield

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{6}n^{(5)}, \frac{1}{3}n^{(5)}, \frac{1}{2}n^{(5)}, \frac{2}{3}n^{(5)}, \frac{5}{6}n^{(5)}, n^{(5)} \right\}.$$

So, it remains to prove that $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{6}n^{(5)}}$, but the proof is the same as in the previous case.

- $n \equiv 2, 10, 14, 18, 22, 30, 34 \pmod{36}$. Using Lemma 2.3 (b) we clearly have for $i = 0, 1, 2, 3, 4$:

- * If $n \equiv 2, 14, 18, 30 \pmod{36}$, then $L_{n+i} \mid F_{\frac{1}{6}n^{(5)}}$.
- * If $n \equiv 10, 22, 34 \pmod{36}$, then $L_{n+i} \mid F_{\frac{1}{2}n^{(5)}}$.

It can be seen that:

- * If $n \equiv 2, 14 \pmod{36}$, then $\gcd\left(\frac{L_n}{3}, L_{n+3}\right) = \gcd\left(\frac{L_n}{3}, \frac{L_{n+4}}{2}\right) = \gcd\left(L_{n+1}, \frac{L_{n+4}}{2}\right) = 1$.
- * If $n \equiv 10 \pmod{36}$, then $\gcd\left(\frac{L_n}{3}, L_{n+3}\right) = \gcd\left(\frac{L_n}{3}, L_{n+4}\right) = \gcd(L_{n+1}, L_{n+4}) = 1$.
- * If $n \equiv 18, 30 \pmod{36}$, then $\gcd\left(\frac{L_n}{2}, L_{n+3}\right) = \gcd\left(\frac{L_n}{2}, \frac{L_{n+4}}{3}\right) = \gcd\left(L_{n+1}, \frac{L_{n+4}}{3}\right) = 1$.
- * If $n \equiv 22, 34 \pmod{36}$, then $\gcd(L_n, L_{n+3}) = \gcd\left(L_n, \frac{L_{n+4}}{3}\right) = \gcd\left(L_{n+1}, \frac{L_{n+4}}{3}\right) = 1$.

By Lemma 2.5 (a), (d) we obtain:

- * If $n \equiv 2, 14, 18, 30 \pmod{36}$, then $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 6F_{\frac{1}{6}n^{(5)}} \mid F_{6\frac{1}{6}n^{(5)}}$.
- * If $n \equiv 10, 22, 34 \pmod{36}$, then $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid 3F_{\frac{1}{2}n^{(5)}} \mid F_{2\frac{1}{2}n^{(5)}}$.

Thus in all cases we have

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \mid n^{(5)}. \quad (3.24)$$

On the other hand, for $i=0, 1, 2, 3, 4$ clearly $L_{n+i} \mid L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}$, hence $2(n+i) \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$. Since:

- * $n \equiv 2, 14 \pmod{36}$, then $n/2, (n+1)/3, n+2, n+3, (n+4)/2$ are pairwise coprime.
- * $n \equiv 10, 22, 34 \pmod{36}$, then $n/2, n+1, n+2, n+3, (n+4)/2$ are pairwise coprime.
- * $n \equiv 18, 30 \pmod{36}$, then $n/2, n+1, n+2, (n+3)/3, (n+4)/2$ are pairwise coprime.

Thus:

- * $n \equiv 2, 14, 18, 30 \pmod{36}$ implies $\frac{1}{6}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$.
- * $n \equiv 10, 22, 34 \pmod{36}$ implies $\frac{1}{2}n^{(5)} \mid z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4})$.

Summarizing, we have:

- * $n \equiv 2, 14, 18, 30 \pmod{36}$ implies

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{k}{6}n^{(5)} : k \in \{0, \dots, 5\} \right\},$$

- * $n \equiv 10, 22, 34 \pmod{36}$ implies

$$z(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}) \in \left\{ \frac{1}{2}n^{(5)}, n^{(5)} \right\}.$$

To prove the assertion for this case we must prove that:

- * $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{2}n^{(5)}}$ if $n \equiv 10, 22, 34 \pmod{36}$, and
- * $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \nmid F_{\frac{1}{3}n^{(5)}}$ and $L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{2}n^{(5)}}$ if $n \equiv 2, 14, 18, 30 \pmod{36}$.

First, we shall prove that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \nmid F_{\frac{1}{3}n^{(5)}} \quad \text{for } n \equiv 2, 14, 18, 30 \pmod{36}. \quad (3.25)$$

In fact, if $n \equiv 2, 30 \pmod{36}$, then

$$\nu_3(L_n L_{n+4}) = 3 > 2 = \nu_3(F_{\frac{1}{3}n^{(5)}}).$$

If $n \equiv 14 \pmod{36}$, then

$$\begin{aligned} \nu_3(L_n L_{n+4}) &= \nu_3(n+4) + 2 > \nu_3(n+4) + 1 \\ &= \nu_3(n) + \nu_3(n+4) = \nu_3(F_{\frac{1}{3}n^{(5)}}). \end{aligned}$$

If $n \equiv 18 \pmod{36}$, then

$$\begin{aligned} \nu_3(L_n L_{n+4}) &= \nu_3(n) + 2 > \nu_3(n) + 1 \\ &= \nu_3(n) + \nu_3(n+4) = \nu_3(F_{\frac{1}{3}n^{(5)}}). \end{aligned}$$

In summary, $L_n L_{n+4} \nmid F_{\frac{1}{3}n^{(5)}}$ for $n \equiv 2, 14, 18, 30 \pmod{36}$.

Now, we shall prove that

$$L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4} \mid F_{\frac{1}{2}n^{(5)}}$$

for all $n \equiv 2, 10, 14, 18, 22, 30, 34 \pmod{36}$. For that, we use the same p -adic valuation argument as before. For $p \neq 2$ and $p \neq 5$, we proceed exactly as in the case $n \equiv 0 \pmod{12}$. For the case $p = 2$, we have

$$\begin{aligned} \nu_2(F_{\frac{1}{2}n^{(5)}}) &= \nu_2(n) + \nu_2(n+2) + \nu_2(n+4) + 1 \\ &\geq 5 > 3 \geq \nu_2(L_n L_{n+1} L_{n+2} L_{n+3} L_{n+4}). \end{aligned}$$

Therefore, the proof is complete. □

REFERENCES

- [1] BENJAMIN, A.—QUINN, J.: *The Fibonacci numbers – Exposed more discretely*, Math. Mag. **76** (2003), 182–192.
- [2] HALTON, J. H.: *On the divisibility properties of Fibonacci numbers*, Fibonacci Quart. **4** (1966), 217–240.
- [3] KALMAN, D.—MENA, R.: *The Fibonacci numbers – exposed*, Math. Mag. **76** (2003), 167–181.
- [4] KOSHY, T.: *Fibonacci and Lucas Numbers with Applications*. Wiley, New York, 2001.
- [5] MARQUES, D.: *On integer numbers with locally smallest order of appearance in the Fibonacci sequence*, Internat. J. Math. Math. Sci., Article ID 407643 (2011), 4 pages.

ORDER OF APPEARANCE OF PRODUCT OF FIVE CONSECUTIVE LUCAS NUMBERS

- [6] MARQUES, D.: *On the order of appearance of integers at most one away from Fibonacci numbers*, Fibonacci Quart. **50** (2012), 36–43.
- [7] MARQUES, D.: *The order of appearance of product of consecutive Fibonacci numbers*, Fibonacci Quart. **50** (2012), 132–139.
- [8] MARQUES, D.: *The order of appearance of powers Fibonacci and Lucas numbers*, Fibonacci Quart. **50** (2012), 239–245.
- [9] MARQUES, D.: *The order of appearance of product of consecutive Lucas numbers*, Fibonacci Quart. **50** (2012), 239–245.
- [10] MARQUES, D.: *Sharper upper bounds for the order of appearance in the Fibonacci sequence*, Fibonacci Quart. **51** (2013), 233–238.
- [11] LAGARIAS, J. C.: *The set of primes dividing the Lucas numbers has density $2/3$* , Pacific J. Math. **118** (1985), 449–461.
- [12] LENGYEL, T.: *The order of the Fibonacci and Lucas numbers*, Fibonacci Quart. **33** (1995), 234–239.
- [13] RIBENBOIM, P.: *My Numbers, My Friends: Popular Lectures on Number Theory*. Springer-Verlag, New York, 2000.
- [14] ROBINSON, D. W.: *The Fibonacci matrix modulo m* , Fibonacci Quart. **1** (1963), 29–36.
- [15] SALLÉ, H. J. A.: *Maximum value for the rank of apparition of integers in recursive sequences*, Fibonacci Quart. **13** (1975), 159–161.
- [16] VINSON, J.: *The relation of the period modulo m to the rank of apparition of m in the Fibonacci sequence*, Fibonacci Quart. **1** (1963), 37–45.
- [17] VOROBIEV, N. N.: *Fibonacci Numbers*. Birkhäuser, Basel, 2003.

Received September 2, 2014

Diego Marques
Departamento de Matemática
Universidade de Brasília
70910-900 Brasília, DF
BRAZIL
E-mail: diego@mat.unb.br

Pavel Trojovský
Department of Mathematics
Faculty of Science
University of Hradec Králové
Rokitanského 62
500 03 Hradec Králové
CZECH REPUBLIC
E-mail: Pavel.Trojovsky@uhk.cz