

INVERSION FORMULAE FOR THE INTEGRAL TRANSFORM ON A LOCALLY COMPACT ZERO-DIMENSIONAL GROUP

FRANCESCO TULONE

ABSTRACT. Generalized inversion formulae for multiplicative integral transform with a kernel defined by characters of a locally compact zero-dimensional abelian group are obtained using a Kurzweil-Henstock type integral.

1. Introduction

In this paper, we consider integral transforms with kernels defined by characters of a locally compact zero-dimensional abelian group. Transforms of this kind are usually called multiplicative transforms (see [1]).

The problem of getting an inversion formula for integral transform is a continual analogue of the one of recovering the coefficients of a convergent series with respect to characters of a compact zero-dimensional abelian group considered in [6]. So, we use those results on series to obtain the correspondent results on transforms. As in [5] and [6], we will use here the Kurzweil-Henstock method of integration.

Our first result is a locally compact version of [6, Theorem 4.2]. The second result is a generalization of [5, Theorem 5.1], where we are weakening the conditions on the way of convergence of the transform.

In comparison with [4], we consider here transforms directly on the group instead of using a mapping of this group on the real line. That mapping was connected with introduction of a certain ordering in this group which can be avoided here.

2000 Mathematics Subject Classification: 26A39, 42C10, 43A25, 43A70.

Keywords: locally compact zero-dimensional abelian group, characters of a group, Kurzweil-Henstock integral, Fourier series, multiplicative integral transform, inversion formula.

2. Preliminaries

Let G be a zero-dimensional locally compact abelian group which satisfies the second countability axiom. We also suppose that the group G is periodic. It is known (see [1]) that a topology in such a group can be given by a chain of subgroups

$$\dots \supset G_{-n} \supset \dots \supset G_{-2} \supset G_{-1} \supset G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots \quad (1)$$

with $G = \bigcup_{n=-\infty}^{+\infty} G_n$ and $\{0\} = \bigcap_{n=-\infty}^{+\infty} G_n$. The subgroups G_n are clopen sets with respect to this topology. As G is periodic, the factor group G_n/G_{n+1} is finite for each n and this implies that G_n (and so, also all its cosets) is compact. Note that the factor group G_n/G_0 is also finite for any $n < 0$, and so, the factor group G/G_0 is countable. We will use the notation given in [6], in particular, we denote by K_n any coset of the subgroup G_n and by $K_n(g)$ the coset of the subgroup G_n which contains the element g . For each $g \in G$ the sequence $\{K_n(g)\}$ is decreasing and $\{g\} = \bigcap_n K_n(g)$.

Now, for each coset K_n of G_n , we choose and fix an element g_{K_n} for the rest of the paper. Then, for each $n \in \mathbb{Z}$, we can represent any element $g \in G$ in the form

$$g = g_{K_n} + \{g\}_n, \quad (2)$$

where $\{g\}_n \in G_n$. Indeed let $g \in K_n$, for some n , then $g = g_{K_n} + g - g_{K_n}$ and we can put $\{g\}_n := g - g_{K_n}$. We agree to put $g_{G_n} = 0$, so that $g = \{g\}_n$ if $g \in G_n$.

Let Γ denotes the dual group of G , i.e., the group of characters of the group G . It is known (see [1]) that under the assumption imposed on G , the group Γ is also a periodic locally compact zero-dimensional abelian group (with respect to the pointwise multiplication of characters) and we can represent it as a sum of increasing sequence of subgroups

$$\dots \supset \Gamma_{-n} \supset \dots \supset \Gamma_{-2} \supset \Gamma_{-1} \supset \Gamma_0 \supset \Gamma_1 \supset \Gamma_2 \supset \dots \supset \Gamma_n \supset \dots \quad (3)$$

introducing a topology in Γ . Then $\Gamma = \bigcup_{i=-\infty}^{+\infty} \Gamma_i$ and $\bigcap_{i=-\infty}^{+\infty} \Gamma_i = \{\gamma^{(0)}\}$, where $(g, \gamma^{(0)}) = 1$ for all $g \in G$ (here and below, (g, γ) denote the value of a character γ at a point g). For each $n \in \mathbb{Z}$ the group Γ_{-n} is the annihilator of G_n , i.e.,

$$\Gamma_{-n} = G_n^\perp := \{\gamma \in \Gamma : (g, \gamma) = 1 \text{ for all } g \in G_n\}.$$

The representation (2), the properties of a character and of the annihilator imply

$$(g, \gamma) = (g_{K_n}, \gamma) \cdot (\{g\}_n, \gamma) = (g_{K_n}, \gamma).$$

So, with a fixed element g_{K_n} , the value (g, γ) is constant for all $g \in K_n$.

The factor groups $\Gamma_{-n-1}/\Gamma_{-n} = G_{n+1}^\perp/G_n^\perp$ and G_n/G_{n+1} are isomorphic (see [1]) and so they are of a finite order for each $n \in \mathbb{Z}$. This implies that the group Γ_{-n}/Γ_0 is also finite for any $n > 0$, and Γ/Γ_0 is countable.

Now, as we have done above for the group G , we choose and fix an element $\gamma_J \in J$ for each coset J of Γ_0 . Then, using multiplication as the group operation on the group Γ , we can represent any element $\gamma \in \Gamma$ in the form:

$$\gamma = \gamma_J \cdot \{\gamma\}, \quad (4)$$

where $\{\gamma\} \in \Gamma_0$. We agree to put $\gamma_{\Gamma_0} = \gamma^{(0)}$, so that $\gamma = \{\gamma\}$ if $\gamma \in \Gamma_0$.

We denote by μ_G and μ_Γ the Haar measures on the groups G and Γ , respectively, and normalize them so that $\mu_G(G_0) = \mu_\Gamma(\Gamma_0) = 1$. We can make these measures complete by including all the subsets of the sets of measure zero into the respective class of measurable sets.

3. Results related to the compact case

We remind here some definitions and results given in the papers [5] and [6]. We consider derivation basis \mathcal{B}_G constituted by the family of basis sets

$$\beta_\nu = \{(I, g) : g \in G, I = K_n(g), n \geq \nu(g)\},$$

where ν runs over the set of all integer-valued functions on G .

In the terminology of the derivation basis theory, any coset K_n , $n \in \mathbb{Z}$, can be called \mathcal{B}_G -interval.

This basis has all the usual properties of a general derivation basis (see [2], [7]).

A β_ν -partition is a finite collection π of elements of β_ν , where the distinct elements (I', g') and (I'', g'') in π have I' and I'' disjoint. If L is a \mathcal{B}_G -interval and $\bigcup_{(I, g) \in \pi} I = L$, then π is called β_ν -partition of L .

For each \mathcal{B}_G -interval L and for any $\beta_\nu \in \mathcal{B}_G$ there exists a β_ν -partition of L .

The following Kurzweil-Henstock type integral was defined in [5]:

DEFINITION 3.1. Let L be a \mathcal{B}_G -interval. A complex-valued function f on L is said to be *Kurzweil-Henstock integrable with respect to basis \mathcal{B}_G* (or H_G -integrable) on L , with H_G -integral A , if for every $\varepsilon > 0$, there exists a function $\nu : L \mapsto \mathbb{Z}$ such that for any β_ν -partition π of L we have:

$$\left| \sum_{(I, g) \in \pi} f(g) \mu_G(I) - A \right| < \varepsilon.$$

We denote the integral value A by $(H_G) \int_L f d\mu_G$.

Remark 3.1. We note that all the above definitions depend on the structure of the sequence of subgroups (1). So, if we consider for the group Γ the definitions of the \mathcal{B}_Γ -basis and the H_Γ -integral, then we should use the sequence (3) in our construction.

Remark 3.2. It is easy to check that H_G -integral is invariant under translation given by some element $g \in G$.

We also need the following extension of Definition 3.1 to the case of functions defined only almost everywhere on L .

DEFINITION 3.2. A complex valued function f defined almost everywhere on a \mathcal{B}_G -interval L is said to be H_G -integrable on L , with integral value A , if the function

$$f_1(g) := \begin{cases} f(g), & \text{where } f \text{ is defined,} \\ 0, & \text{otherwise} \end{cases}$$

is H_G -integrable on L to A in the sense of Definition 3.1.

It is clear that a complex-valued function is H_G -integrable if and only if both its real and imaginary parts are H_G -integrable.

If the group G is compact, as it is in [6], the chain (1) is reduced to the one-side sequence

$$G = G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots$$

In this case the H_G -integral is defined on the whole group G . Moreover, the group Γ of characters of the group G is discrete now (see [1]) and can be represented as a sum of increasing chain of finite subgroups

$$\Gamma_0 \subset \Gamma_{-1} \subset \Gamma_{-2} \subset \dots \subset \Gamma_{-n} \subset \dots,$$

where $\Gamma_0 = \{\gamma^{(0)}\}$ with $(g, \gamma^{(0)}) = 1$ for all $g \in G$.

Characters γ constitute a countable orthonormal system on G with respect to the normalized measure μ_G (see [5]) and we can consider a series

$$\sum_{\gamma \in \Gamma} a_\gamma \gamma \tag{5}$$

with respect to this system. The convergence of this series at a point g is defined in [6] as the convergence of its partial sums

$$S_n(g) := \sum_{\gamma \in \Gamma_{-n}} a_\gamma(g, \gamma) \tag{6}$$

when n tends to infinity.

With the series (5) we associate a function F defined on each coset K_n by

$$F(K_n) := \int_{K_n} S_n(g) d\mu_G. \tag{7}$$

The above integral can be understood in the Lebesgue sense. F is known to be an additive function on the family of all \mathcal{B}_G -intervals.

The next theorem is proved in [6].

THEOREM 3.1. *Suppose that the partial sums $S_n(g)$ of a series $\sum_{\gamma \in \Gamma} a_\gamma \gamma$ and a H_G -integrable function f satisfy the inequalities*

$$\liminf_{n \rightarrow \infty} \operatorname{Re} S_n(g) \leq \operatorname{Re} f(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Re} S_n(g), \quad (8)$$

$$\liminf_{n \rightarrow \infty} \operatorname{Im} S_n(g) \leq \operatorname{Im} f(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Im} S_n(g) \quad (9)$$

everywhere on G except on a countable set S , where

$$\liminf_{n \rightarrow \infty} \mu(K^n) \operatorname{Re} S_n(g) \leq 0 \leq \limsup_{n \rightarrow \infty} \mu(K^n) \operatorname{Re} S_n(g), \quad (10)$$

$$\liminf_{n \rightarrow \infty} \mu(K^n) \operatorname{Im} S_n(g) \leq 0 \leq \limsup_{n \rightarrow \infty} \mu(K^n) \operatorname{Im} S_n(g) \quad (11)$$

hold. Then the series $\sum_{\gamma \in \Gamma} a_\gamma \gamma$ is convergent to f a.e. and it is the H_G -Fourier series of f .

Another version of the theorem on the recovering coefficients was proved in [5]:

THEOREM 3.2. *Suppose that the partial sums $S_n(g)$ of a series $\sum_{\gamma \in \Gamma} a_\gamma \gamma$ converge almost everywhere on G to a function f and satisfy the conditions*

$$-\infty < \liminf_{n \rightarrow \infty} \operatorname{Re} S_n(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Re} S_n(g) < +\infty, \quad (12)$$

$$-\infty < \liminf_{n \rightarrow \infty} \operatorname{Im} S_n(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Im} S_n(g) < +\infty \quad (13)$$

everywhere on G except on a countable set S , where

$$S_n(g) = o\left(\frac{1}{\mu_G(K_n(g))}\right) \quad (14)$$

holds. Then f is H_G -integrable in the sense of Definition 3.2 and the series $\sum_{\gamma \in \Gamma} a_\gamma \gamma$ is the H_G -Fourier series of f .

We need also the following theorem (see [5]).

THEOREM 3.3. *The partial sums $S_n(f, g)$ of the H_G -Fourier series of a H_G -integrable on G function f are convergent to f almost everywhere on G .*

4. Inversion formula for transform in the locally compact case

To simplify our notation, in this section we will put $K = K_0$, $[g] := g_K$, $\{g\} := \{g\}_0$, so that representation (2) with $n = 0$ for any element g of some coset K of G_0 can be rewritten in the form $g = [g] + \{g\}$, where $[g]$ is a fixed element of K and $\{g\} \in G_0$. Similarly, we will sometimes use the notation

$[\gamma] := \gamma_J$ to underline duality, so the representation (4) for any element γ of some coset J of Γ_0 can be rewritten in the form $\gamma = [\gamma] \cdot \{\gamma\}$, where $[\gamma]$ is a fixed element of J and $\{\gamma\} \in \Gamma_0$.

Using this notation and the properties of a character γ , we can write

$$(g, \gamma) = (\{g\}, [\gamma]) \cdot ([g], [\gamma]) \cdot (\{g\}, \{\gamma\}) \cdot ([g], \{\gamma\}). \quad (15)$$

Now we observe that:

- 1) $\{g\} \in G_0$ and $\{\gamma\} \in \Gamma_0 = G_0^\perp$. So $(\{g\}, \{\gamma\}) = 1$, and we can eliminate $(\{g\}, \{\gamma\})$ from representation (15) getting

$$(g, \gamma) = (\{g\}, [\gamma]) \cdot ([g], [\gamma]) \cdot ([g], \{\gamma\}). \quad (16)$$

- 2) $[\gamma] \in \Gamma_{-m(\gamma)} = G_{m(\gamma)}^\perp$ where $m(\gamma) \geq 0$ and $[\gamma]|_{G_0}$ is a character of the subgroup G_0 .
- 3) $([g], [\gamma])$ is constant if g belongs to a fixed coset of G_0 and γ belongs to a fixed coset of Γ_0 .
- 4) Using the duality between G and Γ we can state that g represents a character of Γ and, similarly to the property 2), $[g]|_{\Gamma_0}$ is a character of Γ_0 . So $([g], \{\gamma\})$ is a value of this character at the point $\{\gamma\}$.

Therefore, according to (16), if g belongs to a fixed coset of G_0 and γ belongs to a fixed coset of Γ_0 , we can represent (g, γ) , up to a constant multiplier $([g], [\gamma])$, as a product of $(\{g\}, [\gamma])$ considered as a value of the character $[\gamma]$ at $\{g\}$, and $([g], \{\gamma\})$ considered as a value of the character $[g]$ at $\{\gamma\}$.

We now generalize Theorem 3.1 and Theorem 3.2 to a locally compact case. In what follows, all the integrals are either H_G - or H_Γ -integrals. The measure μ_G or μ_Γ , written under the integral sign, will indicate which of the integrals is used.

THEOREM 4.1. *Assume that G is the group described in Section 2, Γ being its dual group. Let $a(\gamma)$ be a locally H_Γ -integrable function and let for some locally H_G -integrable function ϕ the following inequalities hold:*

$$\liminf_{n \rightarrow \infty} \operatorname{Re} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \leq \operatorname{Re} \phi(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Re} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma, \quad (17)$$

$$\liminf_{n \rightarrow \infty} \operatorname{Im} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \leq \operatorname{Im} \phi(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Im} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \quad (18)$$

everywhere on G except on a countable set T , where we have

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mu_G(K_n) \operatorname{Re} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \\ & \leq 0 \leq \limsup_{n \rightarrow \infty} \mu_G(K_n) \operatorname{Re} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma, \end{aligned} \quad (19)$$

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mu_G(K_n) \operatorname{Im} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \\ & \leq 0 \leq \limsup_{n \rightarrow \infty} \mu_G(K_n) \operatorname{Im} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma. \end{aligned} \quad (20)$$

Then the function $a(\gamma)$ can be recovered from ϕ by the following inversion formula:

$$a(\gamma) = \lim_{n \rightarrow \infty} \int_{G_{-n}} \phi(g) \overline{(g, \gamma)} d\mu_G \quad \text{a.e. on } \Gamma.$$

A particular case of the previous theorem is the following

COROLLARY 4.2. *With the same assumptions on G , Γ and $\alpha(\gamma)$, let*

$$\lim_{n \rightarrow \infty} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma = \phi(g)$$

a.e. on G , where ϕ is a locally H_G -integrable function on G . Moreover, everywhere on G except on a countable set T , we have

$$\limsup_{n \rightarrow \infty} \left| \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \right| < +\infty$$

and for $g \in T$, we have

$$\lim_{n \rightarrow \infty} \mu(K_n(g)) \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu = 0.$$

Then the function $a(\gamma)$ can be recovered from ϕ by the following inversion formula:

$$a(\gamma) = \lim_{n \rightarrow \infty} \int_{G_{-n}} \phi(g) \overline{(g, \gamma)} d\mu_G \quad \text{a.e. on } \Gamma.$$

THEOREM 4.3. *Assume that G is a group described above, Γ being its dual group. Let $a(\gamma)$ be a locally H_Γ -integrable function and*

$$\lim_{n \rightarrow \infty} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma = \phi(g) \quad (21)$$

a.e. on G . Moreover, everywhere on G except on a countable set T , we have

$$\limsup_{n \rightarrow \infty} \left| \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma \right| < +\infty \quad (22)$$

and for $g \in T$ we have

$$\lim_{n \rightarrow \infty} \mu_G(K_n(g)) \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma = 0. \quad (23)$$

Then ϕ is locally H_G -integrable and the function $a(\gamma)$ can be recovered from ϕ by the following inversion formula:

$$a(\gamma) = \lim_{n \rightarrow \infty} \int_{G_{-n}} \phi(g) \overline{\phi(g, \gamma)} d\mu_G \quad \text{a.e. on } \Gamma. \quad (24)$$

Proof of Theorem 4.1 and Theorem 4.3. We sketch the proof following the lines of the proof of a similar result given in [5] indicating the step where we use different theorems on recovering the coefficients of series.

Having fixed a coset K , suppose that $g \in K$, and let J denote any coset of Γ_0 . Then, by (16),

$$\begin{aligned} \int_{\Gamma_{-n}} a(\gamma)(g, \gamma) d\mu_\Gamma &= \sum_{J \subset \Gamma_{-n}} \int_J a(\gamma)(\{g\}, [\gamma]) \cdot ([g], [\gamma]) \cdot ([g], \{\gamma\}) d\mu_\Gamma \\ &= \sum_{J \subset \Gamma_{-n}} (\{g\}, \gamma_J) \cdot \int_J a(\gamma)([g], [\gamma]) \cdot ([g], \{\gamma\}) d\mu_\Gamma. \end{aligned} \quad (25)$$

The latter sum can be considered as a partial sum

$$\sum_{J \subset \Gamma_{-n}} b_J^{(K)}(\{g\}, \gamma_J) \quad (26)$$

of the series with respect to the system of characters $\{\gamma_J\}_J$, at the point $\{g\}$, with the coefficients

$$b_J^{(K)} = \int_J a(\gamma)([g], [\gamma]) (g_K, \{\gamma\}) d\mu_\Gamma.$$

Now, we are going to apply Theorem 3.1 or Theorem 3.2 to this series.

In the case of Theorem 4.1, we obtain, according to the inequalities (17), (18), (19), (20) and the equality (25), that for the last series the following inequalities hold

$$\liminf_{n \rightarrow \infty} \operatorname{Re} S_n(g) \leq \operatorname{Re} \phi(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Re} S_n(g), \quad (27)$$

$$\liminf_{n \rightarrow \infty} \operatorname{Im} S_n(g) \leq \operatorname{Im} \phi(g) \leq \limsup_{n \rightarrow \infty} \operatorname{Im} S_n(g) \quad (28)$$

except on a countable set T where (10) and (11) hold.

We now introduce the variable $t = \{g\} \in G_0$ and we can consider the above inequalities to hold on G_0 for the function $f(t) = \phi(g_K + t)$. This means that the partial sums of our series and the function f satisfy on G_0 all the conditions of Theorem 3.1. Applying this theorem we get that the coefficients $b_J^{(K)}$ are the H_G -Fourier coefficients of $f(t)$, with respect to characters γ_J , i.e.,

$$\begin{aligned} b_J^{(K)} &= \int_J a(\gamma)([g], [\gamma])(g_K, \{g\}) d\mu_\Gamma \\ &= \int_{G_0} f(t) \overline{(\{g\}, \gamma_J)} d\mu_G \\ &= \int_K \phi(g) \overline{(\{g\}, \gamma_J)} d\mu_G \end{aligned} \quad (29)$$

(in the last equality we use Remark 3.2)

In the case of Theorem 4.3, we obtain, according to the assumption (21) and the equality (25) that partial sums (26) are convergent almost everywhere on K to a function $\phi(g)$.

Introducing the variable $t = \{g\} \in G_0$ once again, we can consider this series to be convergent almost everywhere on G_0 to the function $f(t) = \phi(g_K + t)$. The partial sums (26) are bounded, according to (22) and (25), except on a countable set $S = \{t \in G_0 : g_K + t \in T\}$, where (23) holds corresponding to the condition (14) applied to $t \in S$.

Therefore, by Theorem 3.2, the function f is H_G -integrable on G_0 , and so, ϕ is H_G -integrable on K and the coefficients $b_J^{(K)}$ are the H_G -Fourier coefficients of $f(t)$, with respect to characters γ_J getting again the equality (29).

Now, we can finish the proof of both theorems. By observation 3), $([g], [\gamma])$ is constant when $g \in K$ and $\gamma \in J$ with $|([g], [\gamma])| = 1$. Hence (29) implies

$$\int_J a(\gamma)(g_K, \{g\}) d\mu_\Gamma = \int_K \phi(g) \overline{([g], [\gamma])} (\{g\}, \gamma_J) d\mu_G. \quad (30)$$

Now, we notice that for each fixed J , the value $\int_J a(\gamma)(g_K, \{g\}) d\mu_\Gamma$ is the H_G -Fourier coefficient, with respect to the character $\overline{g_K}$, of the H_G -integrable

function $a(\gamma) = a([\gamma] \cdot \{\gamma\})$ considered as a function of $\{\gamma\} \in \Gamma_0$. Applying Theorem 3.3 to this H_G -Fourier series, we get

$$\lim_{n \rightarrow \infty} \sum_{K \subset G_{-n}} \int_J a(\gamma)(g_K, \{\gamma\}) d\mu_\Gamma \cdot \overline{(g_K, \{\gamma\})} = a([\gamma] \cdot \{\gamma\}) = a(\gamma)$$

for almost all values of $\{\gamma\}$ on Γ_0 , i.e., a.e. on J . Hence using (30) and then (16), we compute

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sum_{K \subset G_{-n}} \int_J a(\gamma)(g_K, \{\gamma\}) d\mu_\Gamma \cdot \overline{(g_K, \{\gamma\})} \\ &= \lim_{n \rightarrow \infty} \sum_{K \subset G_{-n}} \int_K \phi(g)([g], [\gamma]) (\{g\}, \gamma_J) d\mu_G \cdot \overline{(g_K, \{\gamma\})} \\ &= \lim_{n \rightarrow \infty} \int_{G_{-n}} \phi(g) \overline{(\{g\}, \gamma_J) \cdot (g_K, \{\gamma\}) \cdot ([g], [\gamma])} d\mu_G \\ &= \lim_{n \rightarrow \infty} \int_{G_{-n}} \phi(g) \overline{(g, \gamma)} d\mu_G = a(\gamma) \quad \text{a.e. on } J. \end{aligned}$$

The last equality is true for any J , so we get (24), completing the proof. \square

We remark that Theorems 4.1 is not true if we use Denjoy-Khintchine integrable function ϕ (see [3] for the compact case), but it becomes true with this last integral if we put some additional hypothesis on the group and on the type of convergence (see [8]).

REFERENCES

- [1] AGAEV, G. N.—VILENKIN, N. YA.—DZHARARLI, G. M.—RUBINSHTEJN, A. I.: *Multiplicative System of Functions and Harmonic Analysis on Zero-Dimensional Groups*. “Ehlm”, Baku, 1981. (In Russian)
- [2] OSTASZEWSKI, K. M.: *Henstock Integration in the Plane*, in: Mem. Amer. Math. Soc. Vol. 63, no. 353, AMS, Providence, RI 1986.
- [3] SKVORTSOV, V.—KOROLEVA, M. P.: *Series in multiplicative systems convergent to Denjoy-integrable functions*, Mat. Sb. **186** (1995), 129–150; Engl. transl. in Sb. Math. **186** (1995), 1821–1842.
- [4] SKVORTSOV, V.—TULONE, F.: *Henstock type integral in harmonic analysis on zero-dimensional groups*, J. Math. Anal. Appl. **322** (2006), 621–628.
- [5] SKVORTSOV, V.—TULONE, F.: *Kurzweil-Henstock type integral on zero-dimensional group and some of its applications*, Czechoslovak Math. J. **58** (2008), 1167–1183.
- [6] SKVORTSOV, V.—TULONE, F.: *Kurzweil-Henstock type integral in Fourier analysis on compact zero-dimensional group*, Tatra Mt. Math. Publ. **44** (2009), 41–51.

INVERSION FORMULAE FOR THE INTEGRAL TRANSFORM ON A GROUP

- [7] THOMSON, B. S.: *Derivation bases on the real line. I., II.*, Real Anal. Exchange **8** (1983), 67–207, 278–442.
- [8] TULONE, F.: *Denjoy and \mathcal{P} -path integrals on compact groups in an inversion formula for multiplicative transforms*, Tatra Mt. Math. Publ. **42** (2008), 27–37.

Received November 20, 2008

*Dipartimento di Matematica ed Applicazioni
Università di Palermo
via Archirafi 34
I-90123 Palermo
ITALY
E-mail: tulone@math.unipa.it*