

MARINE CARBONIFEROUS ALGAE FROM METACARBONATES OF THE OCHTINÁ FORMATION (GEMERIC UNIT, WESTERN CARPATHIANS)

BERNARD MAMET¹ and MILAN MIŠÍK²

¹Departement des sciences de la Terre et de l'environnement, Faculté des Sciences, Université libre de Bruxelles, av. F.D. Roosevelt 50, B-1050 Bruxelles, Belgique

²Department of Geology and Paleontology, Faculty of Sciences, J.A.Comenius University, Mlynská dolina G, 842 15 Bratislava, Slovak Republic

(Manuscript received April 19, 2002; accepted in revised form October 3, 2002)

Abstract: An association of eight taxons of marine algae is described from the anchimetamorphosed dolomites of Late Visean–Namurian A age. The algae confirm this determination of age based previously on conodonts. The flora is indicative of latest Visean Zone 16 to Early Serpukhovian Zone 17. The microfossils from the locality Furmanec are somewhat older, indicative of Late Visean (Zone 15 to 16?).

Key words: Carboniferous, Western Carpathians, Gemic Unit, anchimetamorphosed dolomites, Algae.

Geological part

Algae of Carboniferous age have been reported in the Central Western Carpathians only once (Plašienka & Soták 2001). We therefore described the association found in the Ochtná Formation of the Gemic Unit, mainly from the type locality, an abandoned magnesite quarry near Ochtná and the neighbouring locality of Dúbrava (Fig. 1).

At those localities the Ochtná Formation is represented by bedded dolomites, clayey dolomitic shales with magnesite lenses, graphitic shales and sericitic phyllites. A thick layer of magnesite caps them. The fauna of bivalves, brachiopods and corals from Ochtná was described by Ulrich & Bouček (1931); they considered it to be Uralian (Upper Late Carboniferous). Heritsch (1934) reported an abundant association of corals, which he attributed to the Moscovian (Miatschkovian), while noting a similarity to corals from China. Some specimens were sent to Stanley-Smith; in his opinion they are of Upper Visean age. Mišík (1953) also made some determinations. Trilobites from Ochtná were described by Bouček & Příbyl (1958), who attributed them to the Uppermost Namurian–Westphalian.

However the most accurate determination of age is based mainly on conodonts described by Kozur et al. (1976). The lower faunal horizon includes *Paragnathodus nodosus* (Bischof) representing Uppermost Visean. The upper horizon is dated by "*Gnathodus*" *bilineatus bollandensis* (Higgins et Boschauer) as Namurian A. A brief emergence occurred before the deposition of the second horizon as is shown by the existence of a pisolitic dolocrust 10 cm thick between them (Mišík 1998a, Table I: Figs. 1–3). Turek & Prokop (1981) described a new species of goniatite and chiton, which they attributed to Namurian A. Numerous macrofossils were determined by Macko (1992, in an unpublished manuscript). Bryozoa from a nearby locality Jedlovec were described by Žágoršek & Mac-

ko (1994). They point out their resemblance to those from the Russian platform and the locality Nagyvisnyó in Bükk, Hungary (Westphalian B–C). Sitár & Čapo (1999) report a macroflora dated Westphalian B–C from another neighbouring locality Dúbrava where the enclosing matrix is wackestone and biomicrite. In addition to algae their thin sections contain fragments of corals, bivalves, brachiopods, bryozoans, echinoderm plates, sponge spicules, ostracodes, foraminifers *Pseu-*




Fig. 1. Sketch map showing the position of the localities with marine Carboniferous algae.

doendothyra sp., *Tetrataxis conica* Ehrenberg (determined by J. Kalvoda, University of Brno) and *Globochaete* sp.

The lithology and fauna at Dúbrava indicate that sedimentation took place in the shallow waters of a back-reef and was interrupted by a brief emergence represented by an azoic pisolitic dolocrust (Fig. 2.1). The original character of the Ochtiná Formation was affected by anchimetamorphism. Besides metamorphic foliation, the flakes of new-formed muscovitic mica (Fig. 2.2) and aggregates of light-coloured chlorite replacing bioclasts (Fig. 2.3) can be observed. Both diagenetic and anchimetamorphic silicification are present. As a proof of diagenetic silicification is the presence of authigenic quartz grain with relicts of “dashed” structure (coalescence of parallel shear veinlets) which disappeared through recrystallization in another part of the veinlet (Fig. 2.4; more see Mišík 1998b). Anchimetamorphic silicification is responsible for irregular replacement of bioclasts and recrystallized dolomite (pseudodolosparite) by both microquartz and megaquartz (Fig. 2.5). Rare authigenic feldspars with sparse twins formed during the anchimetamorphism (Fig. 2.6), do not have the precise boundaries commonly ascribed to diagenetic feldspars crystallized during the course of the “pre-metamorphic stage”. The pyrite cubes to 0.15 mm belong to the youngest elements.

Paleontological part

The algal flora is somewhat difficult to identify as tectonic reorientation, stress and incipient metamorphism usually destroy most of the original floral structures. There are, however, a few identifiable elements:

Calcifolium cf. *punctatum* Maslov, 1956
Claracrusta catenoides Homann, 1972
 Dissolved Epimastopodid? sp.
Exvotarissella? sp.
Paraepimastopora sp.
Praedonezella cespeformis Kulik, 1973
Ungdarella uralica Maslov, 1956

Calcifolium (two valid species: *okense* Schvetsov et Birina, 1935 and *punctatum* Maslov, 1956) is one of the rare Paleozoic genera to have a rather precise connotation regarding age. In the Tethys, it straddles the Visean/Namurian boundary. *Okense* is abundant both above and below it, *punctatum* is abundant in the latest Visean. The genus is not known in the American flora (Mamet & Roux 1977; Skompski 1981; Skompski et al. 1989; Bogush et al. 1990; Sebbar 1998).

Claracrusta Vachard, 1981, *Berestovia* Berchenko, 1982 and *Iberiaella* Racz, 1984 cause many taxonomical problems, so they are of little value in answering the stratigraphic question posed here. Indeed *Claracrusta* and the type *C. catenoides* (Homann, 1972) range through the Carboniferous and Permian. The oldest *catenoides* are illustrated from the latest Visean (Sebbar & Mamet 1996) and the youngest are Murghabian (Flügel 1990). Vachard & Maslo (1996) mention its presence in the Late Permian, but without convincing illustrations.

Epimastopodid? and *Exvotarissella*? sp. cannot be identified with certitude.

The genus *Paleoberesella* is present as early as the mid-Devonian (Mamet & Pr  at 1992) and its greatest expansion is in the Visean (Mamet & Roux 1974). It ranges into the mid-Carboniferous.

The oldest representative of *Paraepimastopora* are illustrated by Sanchez-Chico et al. (1995) from the Late Visean (Zone 15, Spain) and Sebbar & Mamet (1996) from the Early Serpukhovian (Zone 17, Algeria). It is also known from many other Carboniferous localities. Thus the range reported by Granier & Deloffre (1994) should be considerably extended downward in the Carboniferous.

Praedonezella cespeformis Kulik, 1973, is a Tethyan Late Visean to early Bashkirian taxon (Bogush et al. 1990; Chuvashov et al. 1993).

The latest identifiable taxon, the rhodophyte *Ungdarella uralica* Maslov, 1956 (non 1950) is long ranging, as it is known from the Late Visean Zone 16 (as *Ungdarella decean-glorum* Elliott, 1957 and later on by Vachard et Fadli, 1991). It is illustrated from the Lower Serpukhovian–Early Namurian by Mamet & Roux (1977) and Sebbar (1998).

The flora is therefore composed of 8 identifiable genera, 6 of which are long ranging and of little stratigraphic use. On the other hand *Calcifolium* and *Praedonezella* are more restricted in range. This Tethyan flora is indicative of latest Visean Zone 16, to Early Serpukhovian, Zone 17.

Comparison

Pla  ienka & Sot  k (2001) have recently described Carboniferous microfossils from the locality Furmanec, (Veporic Unit) that contains not only algae (Stacheins and Paleoberesellids), but also foraminifers. They reported an association of *Valvulinella latissima* Conil et Lys, *Millerella concinna* Potievskaya and *Eotextularia diversa* (Chernysheva). The first species is Late Visean, the second genus Bashkirian and the third taxon Early Visean. They should be transferred to *Valvulinella youngi* (Brady), *Eostafella* sp. and *Paleotextularia sensu stricto*. Thus the fauna is indicative of Late Visean (Zone 15? to 16?) an age somewhat older than Ochtin   (G  meric Unit).

Fig. 2. Some microscopic features of algae-bearing dolomites. 1 — Dolocrust with vadose pisolites, indicating a temporary interruption of marine sedimentation. Namurian A, Ochtin   Formation. Abandoned magnesite quarry near Ochtin  , Spi  sko-gemersk   rudohorie Mts. Thin section No. 1718. 2 — Newly-formed micas in the anchimetamorphosed dolomite with intraclasts. Upper Visean–Lower Namurian. As above. T.s. 18931. 3 — Bioclast replaced by chlorite aggregate. As above. T.s. 18927. 4 — Calcite veinlet originated by recrystallization of parallel shear veinlets. Their relict existence is preserved only where replaced by early diagenetic authigenic quartz (white). As above. 5 — Partly silicified algal aggregate and pseudodolosparite. As above. T.s. 18933. 6 — Authigenic feldspars in the anchimetamorphosed dolomite. Upper Visean–Lower Namurian. As above. T.s. 19010.




Fig. 2.









Fig. 3.

Fig. 3. Association of algae. 1 — *Paraepimastopora* sp., cross-section in the dolomite of Upper Viséan–Lower Namurian of Gemeric Unit. Dúbrava near Jelšava. T.s. 24975. 2 — *Paraepimastopora* sp., tangential section. As above. 3 — *Paraepimastopora* sp., oblique section. As above. T.s. 23663. 4 — *Ungdarella* sp., with partially silicified margins. As above. T.s. 24972. 5 — *Calcifolium* cf. *punctatum* Maslov, 1956. Longitudinal section. As above. T.s. 24975. 6 — *Praedonezella cespeformis* Kulik 1973 in Upper Viséan–Lower Namurian dolomites. Its stratigraphic span is from Upper Viséan to Serpukhovian. Abandoned magnesite quarry near Ochtná. T.s. 19003.

Conclusions

Paleozoic algae are often long ranging and far more useful for reconstruction of paleoenvironments than for zonation. This report is therefore exceptional as it provides a rather restricted age for rock which have been extensively altered by anchimetamorphism.

References

- Berchenko O.I. 1982: New species of the Upper Serpukhovian deposits of the Donbass depression. In: Teslenko Yu.V. (Ed.): Systematics and evolution of fossil plants from the Ukraine. *Dumka Naukova*, Kiev, 51–55 (in Russian).
- Bogush O.I., Ivanova R.M. & Luchinina V.A. 1990: Calcareous algae of the Late Famennian and Early Carboniferous from the Urals and Siberia. *Trudy Inst. Geol. Geofiz. Akad. Nauk SSSR, Sibirskoe Otdelenie* 745, 1–160.
- Bouček B. & Přibyl A. 1958: Preliminary note concerning the investigation of marine Carboniferous in the Spiš-Gemer Metaliferous Mts. *Geol. Práce, Spr.* 14, 118–123 (in Czech).
- Bouček B. & Přibyl A. 1960: Revision der Trilobiten aus dem slowakischen Oberkarbon. *Geol. Práce, Spr.* 20, 5–50.
- Chuvashov B.J., Shuysky V.P. & Ivanova R.M. 1993: Stratigraphical and facies complexes of the Paleozoic calcareous algae of the Urals. In: Barattolo et al. (Eds.): Studies of fossil benthic algae. *Boll. Soc. Paleont. Ital. Spec. Vol.* 1, 93–119.
- Elliott G.F. 1970: Calcareous algae new to the British Carboniferous. *Paleontology* 13, 3, 443–450.
- Flügel E. 1966: Algen aus dem Perm der Karnischen Alpen. *Carinthia* II, 25, 3–76.
- Flügel E. 1990: "Einschnitte" in der Entwicklung permischer Kalkalgen. *Mitt. Naturwiss. Ver. Steiermark* 20, 99–124.
- Granier B. & Deloffre R. 1994: Inventaire critique des Algues dasycladales fossiles. IIIe partie. Les Algues dasycladales du Permien et du Trias. *Rev. Paléobiologie* 14, 1, 49–84.
- Heritsch F. 1934: Rugose Korallen aus der Karbon der tschechoslowakischen Karpathen. *Věst. SGÚ* 10, 138–154.
- Homann W. 1972: Unter- und tief-mittelpermische Kalkalgen aus dem Rattendorfer Schichten, dem Trogkofel-Kalk und dem Tressdorfer Kalk der Karnischen Alpen (Oesterreich). *Senckenberg. Lethaea* 53, 133–313.
- Kozur H., Mock R. & Mostler H. 1976: Stratigraphische Neueinstufung der Karbonatgesteine der unteren Schichtenfolge von Ochtná (Slowakei) in das oberste Visé und Serpukhovian (Namur A). *Geol. Paläont. Mitt., Innsbruck* 6, 1, 1–29.
- Kulik E.L. 1973: Cyanophyta. In: Einor O.L. (Ed.): Stratigraphy and fauna of the Carboniferous succession of the Shartym River (southern Urals). *Uralskoe geologicheskoe Upravlenie, Izd. Lvov. Gosud. Universiteta*, Lvov, 1–175.
- Macko A. 1992: Geology and stratigraphy of marine Carboniferous in Ochtná. *Dissertation. Manuscript (Archive of Department of Geology and Paleontology, J. A. Comenius University, Bratislava)*, 1–37 (in Slovak).
- Malik K. 1978: Sedimentological study of Grodzisce (Hradište) Conglomerates and microfacies of their exotic carbonate pebbles. *Manuscript. Archive Inst. Nauk Geol.*, Kraków, 1–218 (in Polish).
- Mamet B. & Prétat A. 1982: Algues du Dévonien Moyen de Wellin (Synclorium de Dinant, Belgique). *Rev. Micropaléont.* 35, 1, 53–75.
- Mamet B. & Roux A. 1974: Sur quelques Algues tubulaires scalariformes de la Téthys paléozoïque. *Rev. Micropaléont.* 17, 3, 134–156.
- Mamet B. & Roux A. 1975: Algues dévoniennes et carbonifères de la Téthys occidentale. *Rev. Micropaléont.* 17, 3, 134–187.
- Mamet B. & Roux A. 1977: Algues rouges dévoniennes et carbonifères de la Téthys occidentale. *Rev. Micropaléont.* 19, 4, 215–266.
- Maslov V.P. 1956: Fossil calcareous algae of the USSR. *Trudy Inst. Geol. Nauk SSSR* 160, 1–301.
- Mišík M. 1953: Geological investigation of the area between Jelšava and Štítnik. *Geol. Sbor. SAV* 4, 3–4, 557–587 (in Slovak).
- Mišík M. 1994: Authigenic feldspars in carbonate rocks of the Western Carpathians. *Geol. Carpathica* 45, 2, 103–111.
- Mišík M. 1998a: Stratigraphical horizons and facies with calcareous oncoids, microoncoids and pisoids in Western Carpathians. *Miner. Slovaca* 30, 3, 195–216 (in Slovak with English summary).
- Mišík M. 1998b: Peculiar types of thin veins in the Mesozoic carbonates and silicites of the Western Carpathians. *Geol. Carpathica* 49, 4, 271–287.
- Plašienka D. & Soták J. 2001: Stratigraphic and tectonic position of Carboniferous sediments in the Furmanec (Murán Plateau, Central Western Carpathians). *Miner. Slovaca* 33, 1, 29–44 (in Slovak with English summary).
- Racz L.G. 1984: *Iberiella*, a new fossil alga from the Middle Carboniferous of NW Spain. *Geol. en Mijnbouw* 63, 333–336.
- Sanchez-Chico F., Mamet B., Moreno-Eiris E. & Rodriguez S. 1995: Algas calcáreas del Viséense de los Santos de Maimona. *Rev. Esp. Micropaleontol.* 27, 3, 67–96.
- Schvetsov D.V. & Birina L.M. 1935: On the origin and petrology of the Oka Limestone. *Trudy Mosk. Geol. Tresta* 10, 20–21.
- Sebbar A. 1998: Foraminifères et Algues calcaires du Carbonifère, bassin de Reggane, Sahara central, Algérie. *Bull. Serv. Géol. d'Algérie* 9, 2, 123–147.
- Sebbar A. & Mamet B. 1996: Algues benthiques calcaires du Carbonifère Inférieur et Moyen, bassin de Béchar, Algérie. *Rev. Micropaléont.* 39, 2, 153–167.
- Sitár V. & Čapo J. 1999: Carboniferous flora of the Dúbrava masif magnesite deposit. *Miner. Slovaca* 31, 5–6, 525–528 (in Slovak).
- Skompski S. 1981: Morphology and systematic position of the Carboniferous algal genus *Calcifolium*. *Neu. Jb. Geol. Paläont., Mh.* 3, 165–179.
- Skompski S., Conil R., Laloux M. & Lys M. 1989: Etude micropaléontologique des calcaires du Viséen terminal et du Namurien dans le bassin de Lublin à l'Est de la Pologne. *Bull. Soc. Belge Géol.* 98, 3/4, 353–69.
- Turek V. & Prokop R. J. 1982: Two remarkable finds of molluscs in the Carboniferous of Slovakia. *Čas. Mineral. Geol. Praha* 27, 3, 285–294.
- Ulrich F. & Bouček B. 1931: Occurrences of fossils in the magnes-

- ite zone of Slovak Metalliferous Mountains. *Věst. SGÚ* 7 (in Czech).
- Vachard D. 1981: Téthys et Gondwana au Paléozoïque supérieur. Les données afghanes. *Documents et Travaux. Inst. Géol. A. de L'apparent (IGAL)*, Paris 2, 1–463.
- Vachard D. & Maslo A. 1966: Précisions biostratigraphiques et micropaléontologiques sur le Bashkirien d'Ukraine. *Rev. Paléobiologie* 15/2, 357–383.
- Zágoršek K. & Macko A. 1994: Carboniferous Bryozoa from the Jedlovec quarry in Ochtiná Formation (Gemericum, Western Carpathians). *Miner. Slovaca* 26, 5, 335–346 (in Slovak with English summary).