

## DISTRIBUTION OF MOUSE CYTOMEGALOVIRUS IN ORGANS OF WHITE MICE EXPERIMENTALLY INFECTED BY NATURAL ROUTE

J. SVOBODOVÁ, \*D. BLÁŠKOVIČ, D. HUČKOVÁ

Department of Virology and Microbiology, Faculty of Natural Sciences, Comenius University and \*Institute of Virology, Slovak Academy of Sciences, 817 03 Bratislava, Czechoslovakia

Received May 22, 1986

*Summary.* — One, 10, 21-day-old and adult mice were inoculated by peroral and/or intranasal routes with mouse cytomegalovirus (MCMV). In animals surviving generalized infection, the virus could be demonstrated in salivary glands up to 123 days post-infection (p. i.). In mouse females which had eaten their infected and diseased offspring, the virus was detectable in salivary glands up to day 121, p. i. On day 16 p. i., the virus was present in salivary glands, lungs and kidneys of mice of different age groups, but no virus was recovered from their Gasserian ganglia. These results were compared with those obtained after infection with murine alphaherpesvirus.

*Key words:* mouse cytomegalovirus; experimental infection; virus distribution

### Introduction

Pathogenetic studies with murine herpesvirus (MHV) (Blaškovič *et al.*, 1980, Čiampor *et al.*, 1981; Svobodová *et al.*, 1982a, b) in mice infected by natural route demonstrated the presence of this virus in various organs of infected animals and the dynamic nature of its persistence in lungs and latency in Gasserian ganglia (Blaškovič *et al.*, 1984; Rajčáni *et al.*, 1985). No virus was demonstrated in the salivary glands of MHV-infected laboratory mice. Because the population of free living small rodents *Apodemus flavicollis* and *Clethrionomys glareolus* in Czechoslovakia (Mistříková and Blaškovič, 1985) showed positivity for both MHV and MCMV, it appeared reasonable to compare the persistence of MCMV in salivary glands with the persistence of MHV in Gasserian ganglia. This study was performed also with the aim to find further biological criteria distinguishing between MHV (subfamily Alphaherpesvirinae) (Matthews, 1982) and MCMV.

### Materials and Methods

Outbred 1, 10, 21 day old and adult white mice from Dobrá Voda breed were used throughout. In addition, females which have eaten their infected and diseased offspring were included into the experiments. The route of infection and the infectious dose of MCMV strains Smith (1954)

**Table 1. One-day-old suckling mice infected by oral route with  $10^{3.5}$  TCID<sub>50</sub> in 0.01 ml of MCMV (strain Smith)**

Organs	Days p.i.							
	2	4	5	7	10	12	13	16
Salivary glands	+	+	+	+	+	+	+	+
Lungs	+	+	+	+	+	+	+	+
Liver	—	—	n.d.	n.d.	n.d.	n.d.	—	—
Spleen	+	—	+	+	tox	n.d.	+	+
Kidney	+	+	+	+	+	+	+	+
Heart	+	+	+	+	+	n.d.	—	—
Brain	—	—	n.d.	n.d.	n.d.	n.d.	—	—
Gasserian ganglia	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	—	—

n.d. = not done; — = no virus recovered; tox = toxic suspension

used in a titre of  $10^{4.5}$  TCID<sub>50</sub>/0.1 ml imitated natural infection. Volumes of 0.01, 0.02, 0.05 and 0.1 ml respectively, were given perorally (p.o.) and/or intranasally (i.n.).

The presence of the virus distributed via blood to different organs of infected animals was proved by inoculation of the 10% homogenized organ suspension to the primary mouse embryo cell (MEC) monolayer cultures cultivated under standard conditions in basal Eagle's medium supplemented with 10 per cent inactivated bovine serum.

### Results

The virus distribution at different time intervals p. i. in one-day-old mice is presented in Table 1. The infected suckling mice became ill and the severe diseased offspring were eaten by their mothers. When these were investigated on days 76, 89 and 121, MCMV was recovered from their salivary glands, but not from their lungs, spleen and kidneys. Ten days old mice have not developed any overt symptoms of disease. Animals killed by aether

**Table 2. Ten day-old suckling mice infected by oral route with  $2 \times 10^{3.5}$  TCID<sub>50</sub> in 0.02 ml of MCMV**

Organs	Days p.i.				
	2	5	8	13	16
Salivary glands	+	+	n.d.	+	+
Lungs	+	+	+	+	+
Liver	—	—	n.d.	—	—
Spleen	+	+	n.d.	—	—
Kidney	+	+	n.d.	—	+
Brain	n.d.	—	n.d.	—	n.d.
Gasserian ganglia	n.d.	—	n.d.	—	n.d.
Tonsils, mouth mucosa	+	+	n.d.	n.d.	n.d.
Trachea	n.d.	+	n.d.	n.d.	n.d.

Explanation as in Table 1.

Table 3. 21-day-old mice infected by oral route with  $5 \times 10^{3.5}$  TCID<sub>50</sub> in 0.05 ml of MCMV

Organs	Days p.i.				
	3	5	7	10	18
Salivary glands	+	+	+	+	+
Lungs	+	+	+	—	—
Liver	—	—	—	—	—
Spleen	+	+	+	—	—
Kidney	—	+	+	—	n.d.
Heart	+	—	—	—	—
Brain	—	—	—	—	—
Gasserian ganglia	n.d.	—	—	—	—

Explanation as in Table 1.

anaesthesia at different time intervals revealed the presence of infectious virus in organ suspensions (Table 2).

In 21-day-old mice inoculated similarly as the 10-day-old animals the virus could be detected up to day 18 as indicated in Table 3. Adult mice infected with 0.1 ml of virus with infectious titre of  $10^{4.5}$  TCID<sub>50</sub>/0.1 ml were followed for virus persistence. Salivary glands, lungs, spleen and kidneys were investigated on days 31, 42 and 123 p. i. At these intervals the virus was constantly present in salivary glands, in lungs up to the day 31 p. i., its titre in organs reaching  $10^{5.5}$  TCID<sub>50</sub>/ml. No virus was recovered from spleen and kidneys at any of these intervals.

### Discussion

The aim of this study was to compare the pathogenesis of MHV (Alpha-herpesvirinae) first isolated in our laboratory in 1980 with MCMV (Beta-herpesvirinae). The pathogenesis of MHV after experimental infection resembling natural conditions of virus infection and its possible transmission (horizontal infection from mother to the offspring, excretion by nasal or oral discharge and by urine to the environment) were previously described (Blaškovič *et al.*, 1984; Rajčani *et al.*, 1985). The pathogenesis of MCMV was described by Lussier (1973), Hudson (1979) and Tonari and Minamishima (1983). The main interest was paid to long-term virus persistence in different organs of experimental animals to investigate its significance in nature and possibly also in human infections with the viruses of two different genera. MHV persisted in lungs, kidneys and brain of 1-day-old mice for at least 9 days p. i. Ten-day-old animals harboured the virus for at least 11 days in Gasserian ganglia, 14 days in kidneys and 141 days in lungs. In 21 day old mice the MHV could be recovered from Gasserian gnaglia and from kidney up to day 28 and from the lungs up to day 141 p. i. The horizontal transmission from suckling mice to their mothers (oral infection) was effective at least 141–169 days after the presumed infection and the virus could be recovered from Gasserian ganglia and lungs. The virus spread by blood



stream; the persistence in the lungs was of productive type in contrast to the static, nonproductive persistence of herpes simplex virus in experimentally infected rabbits and mice (Rajčáni *et al.*, 1969; Baringer, 1975; Stevens, 1975; Klein, 1982; Rajčáni and Blaškovič, 1986).

In comparison to MHV, the MCMV also spreads by blood stream. The persistence of the virus in salivary gland was decisive for the pathogenesis and for virus excretion. It was accompanied with short-term virus persistence in kidneys and possible virus excretion in urine. No virus presence was detected in neural tissue. The site of virus persistence in lungs and sensory ganglia (MHV), or in salivary glands and lungs (MCMV) and their character may serve as biological marker for distinction of herpesviruses of subfamily Alpha- and Betaherpesvirinae.

*Acknowledgement.* The authors are indebted to Dr. J. Rajčáni for his valuable suggestions and to Mrs. E. Trenčianska for her skilful technical assistance.

#### References

- Beringer, J. R. (1975): Herpes simplex virus infection of nervous tissue in animals and man. *Progr. med. Virol.* **20**, 1–16.
- Blaškovič, D., Stančková, M., Svobodová, J., and Mistríková, J. (1980): Isolation of five strains of herpesviruses from two species of free living small rodents. *Acta virol.* **24**, 468.
- Blaškovič, D., Stančková, D., and Rajčáni, J. (1984): Experimental pathogenesis of murine herpesvirus in newborn mice. *Acta virol.* **28**, 225–231.
- Čiampor, F., Stančková, M., and Blaškovič, D. (1981): Electron microscopy of rabbit embryo fibroblasts infected with herpesvirus isolated from *Clethrionomys glareolus* and *Apodemus flavicollis*. *Acta virol.* **25**, 101–107.
- Hudson, J. B. (1979): The murine cytomegalovirus as a model for study of viral pathogenesis and persistent infections. *Arch. Virol.* **62**, 1–29.
- Klein, J. R. (1982): Pathogenesis of acute, latent and recurrent herpes simplex virus infections. *Arch. Virol.* **72**, 143–168.
- Lussier, G. (1973): Encephalitis caused by murine cytomegalovirus in newborn and weanling mice. *Vet. Path.* **10**, 366–374.
- Matthews, R. E. F. (1982): *Classification and Nomenclature of Viruses*, p. 47–51. Karger, Basel.
- Mistríková, J., and Blaškovič, D. (1985): Ecology of the murine alphaherpesvirus and its isolation from lungs of rodents in cell culture. *Acta virol.* **29**, 312–317.
- Rajčáni, J., Sabó, A., and Blaškovič, D. (1969): Vergleichende Untersuchungen zur experimentellen Pathogenese des Herpes simplex Virus bei der Maus. Intracerebrale, intraperitoneale und orale Infektion des Säuglingmaus. *Zentbl. Bakt. ParasitKde Abt. I* **215**, 1–15.
- Rajčáni, J., Blaškovič, D., Svobodová, J., Čiampor, F., Hučková, D., and Stančková, D. (1985): Pathogenesis of acute and persistent murine herpesvirus infection in mice. *Acta virol.* **29**, 51–60.
- Rajčáni, J., and Blaškovič, D. (1986): Comparative studies on experimental latent herpesvirus infections. *Rev. roum. Virol.*, in press.
- Smith, M. G. (1954): Propagation of salivary gland virus of the mouse in tissue cultures. *Proc. Soc. exp. Biol. Med.* **86**, 435–440.
- Stevens, J. G. (1975): Latent herpes simplex virus and nervous system. *Curr. Top. Microbiol. Immunol.* **70**, 31–50.
- Svobodová, J., Blaškovič, D., and Mistríková, J. (1982a): Growth characteristics of herpesviruses isolated from free living small rodents. *Acta virol.* **26**, 256–263.
- Svobodová, J., Stančková, M., Blaškovič, D., Mistríková, J., Leššo, J., Russ, G., and Masárová, P. (1982b): Antigenic relatedness of alphaherpesviruses isolated from free living rodents. *Acta virol.* **26**, 438–443.
- Tonari, Y., and Minamishima, Y. (1983): Pathogenicity and immunogenicity of temperature-sensitive mutants of murine cytomegalovirus. *J. gen. Virol.* **64**, 1983–1990.