

INEQUALITIES FOR POWER SERIES WITH NONNEGATIVE COEFFICIENTS VIA REVERSES OF JENSEN INEQUALITY

SEVER S. DRAGOMIR

ABSTRACT. Some inequalities for power series with nonnegative coefficients via a reverse of Jensen inequality obtained by Dragomir and Ionescu in 1994 are given. Applications for some fundamental functions defined by power series are also provided.

1. Introduction

In 1994, S. S. Dragomir and N. M. Ionescu obtained the following reverse of Jensen’s discrete inequality:

Let $\Phi : I \rightarrow \mathbb{R}$ be a differentiable convex function on the interior $\overset{\circ}{I}$ of the interval I . If $x_i \in \overset{\circ}{I}$ and $w_i \geq 0$ ($i = 1, \dots, n$) with $W_n := \sum_{i=1}^n w_i = 1$, then one has the inequality:

$$0 \leq \sum_{i=1}^n w_i \Phi(x_i) - \Phi\left(\sum_{i=1}^n w_i x_i\right) \leq \sum_{i=1}^n w_i \Phi'(x_i) x_i - \sum_{i=1}^n w_i \Phi'(x_i) \sum_{i=1}^n w_i x_i. \quad (1.1)$$

In order to improve Grüss’ discrete inequality, P. Cerone and S. S. Dragomir established in 2002 the following result [1]:

$$\left| \sum_{i=1}^n w_i a_i b_i - \sum_{i=1}^n w_i a_i \sum_{i=1}^n w_i b_i \right| \leq \frac{1}{2} (A - a) \sum_{i=1}^n w_i \left| b_i - \sum_{j=1}^n w_j b_j \right|$$

$$\leq \frac{1}{2} (A - a) \left[\sum_{i=1}^n w_i b_i^2 - \left(\sum_{i=1}^n w_i b_i \right)^2 \right]^{1/2}, \quad (1.2)$$

provided $-\infty < a \leq a_i \leq A < \infty$, and $w_i \geq 0$ ($i = 1, \dots, n$) with $W_n := \sum_{i=1}^n w_i = 1$.

© 2016 Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: 26D15, 26D10.

Keywords: power series, Jensen’s inequality, reverse of Jensen’s inequality.

In addition, if $-\infty < b \leq b_i \leq B < \infty$, ($i = 1, \dots, n$), then we have the string of inequalities

$$\begin{aligned} & \left| \sum_{i=1}^n w_i a_i b_i - \sum_{i=1}^n w_i a_i \sum_{i=1}^n w_i b_i \right| \\ & \leq \frac{1}{2} (A - a) \left[\sum_{i=1}^n w_i b_i^2 - \left(\sum_{i=1}^n w_i b_i \right)^2 \right]^{1/2} \leq \frac{1}{4} (A - a) (B - b). \end{aligned} \quad (1.3)$$

Utilising these results, we observe that if Φ is differentiable convex on a finite interval, say $[m, M]$, then we have the inequalities:

$$\begin{aligned} 0 & \leq \sum_{i=1}^n w_i \Phi(x_i) - \Phi \left(\sum_{i=1}^n w_i x_i \right) \\ & \leq \sum_{i=1}^n w_i \Phi'(x_i) x_i - \sum_{i=1}^n w_i \Phi'(x_i) \sum_{i=1}^n w_i x_i \\ & \leq \frac{1}{2} (M - m) \sum_{i=1}^n w_i \left| \Phi'(x_i) - \sum_{j=1}^n w_j \Phi'(x_j) \right| \\ & \leq \frac{1}{2} (M - m) \left[\sum_{i=1}^n w_i [\Phi'(x_i)]^2 - \left(\sum_{i=1}^n w_i \Phi'(x_i) \right)^2 \right]^{1/2} \end{aligned} \quad (1.4)$$

for $x_i \in (m, M)$ ($i = 1, \dots, n$).

If the lateral derivatives $\Phi'_+(m)$ and $\Phi'_-(M)$ are finite, then we also have

$$\begin{aligned} 0 & \leq \sum_{i=1}^n w_i \Phi(x_i) - \Phi \left(\sum_{i=1}^n w_i x_i \right) \\ & \leq \frac{1}{2} [\Phi'_-(M) - \Phi'_+(m)] \sum_{i=1}^n w_i \left| x_i - \sum_{j=1}^n w_j x_j \right| \\ & \leq \frac{1}{2} [\Phi'_-(M) - \Phi'_+(m)] \left[\sum_{i=1}^n w_i x_i^2 - \left(\sum_{i=1}^n w_i x_i \right)^2 \right]^{1/2} \\ & \leq \frac{1}{4} (M - m) [\Phi'_-(M) - \Phi'_+(m)] \end{aligned} \quad (1.5)$$

for $x_i \in [m, M]$ ($i = 1, \dots, n$).

In the recent paper [9], by the use of a refinement of Young's inequality, the authors proved the following result:

Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $y, z, z^\nu y^{1-\nu}, z^{1-\nu} y^\nu \in (0, R)$ and $\nu \in [0, 1]$, then we have the inequalities:

$$\begin{aligned} & 2 \min \{ \nu, 1 - \nu \} [f(y) f(z) - f^2(\sqrt{yz})] \\ & \leq f(y) f(z) - f(z^\nu y^{1-\nu}) f(z^{1-\nu} y^\nu) \\ & \leq 2 \max \{ \nu, 1 - \nu \} [f(y) f(z) - f^2(\sqrt{yz})] \end{aligned} \tag{1.6}$$

or, equivalently,

$$\begin{aligned} & 2 \min \{ \nu, 1 - \nu \} [f(u^2) f(t^2) - f^2(ut)] \\ & \leq f(u^2) f(t^2) - f(u^{2\nu} t^{2(1-\nu)}) f(t^{2(1-\nu)} u^{2\nu}) \\ & \leq 2 \max \{ \nu, 1 - \nu \} [f(u^2) f(t^2) - f^2(ut)], \end{aligned} \tag{1.7}$$

provided $u^2, t^2, u^{2\nu} t^{2(1-\nu)}, t^{2(1-\nu)} u^{2\nu} \in (0, R)$ and $\nu \in [0, 1]$.

For other recent results for power series with nonnegative coefficients, see [9] and [10]. For more results on power series inequalities, see [2] and [5]–[8].

Motivated by the above results and utilizing a reverse of Jensen inequality obtained by S. S. Dragomir and N. M. Ionescu in 1994, we provide in this paper other inequalities for power series with nonnegative coefficients. Applications for some fundamental functions are given as well.

2. Power inequalities

The most important power series with nonnegative coefficients are:

$$\begin{aligned} \exp(z) &= \sum_{n=0}^{\infty} \frac{1}{n!} z^n, & z \in \mathbb{C}, \\ \frac{1}{1-z} &= \sum_{n=0}^{\infty} z^n, & z \in D(0, 1), \\ \ln \frac{1}{1-z} &= \sum_{n=1}^{\infty} \frac{1}{n} z^n, & z \in D(0, 1), \\ \cosh z &= \sum_{n=0}^{\infty} \frac{1}{(2n)!} z^{2n}, & z \in \mathbb{C}, \\ \sinh z &= \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} z^{2n+1}, & z \in \mathbb{C}. \end{aligned} \tag{2.1}$$

Other important examples of functions as power series representations with non-negative coefficients are:

$$\begin{aligned} \frac{1}{2} \ln \left(\frac{1+z}{1-z} \right) &= \tanh^{-1}(z) = \sum_{n=1}^{\infty} \frac{1}{2n-1} z^{2n-1}, & z \in D(0,1), \\ \sin^{-1}(z) &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\frac{1}{2})}{\sqrt{\pi}(2n+1)n!} z^{2n+1}, & z \in D(0,1), \end{aligned} \tag{2.2}$$

$${}_2F_1(\alpha, \beta, \gamma, z) := \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)\Gamma(n+\beta)\Gamma(\gamma)}{n!\Gamma(\alpha)\Gamma(\beta)\Gamma(n+\gamma)} z^n, \alpha, \beta, \gamma > 0, \quad z \in D(0,1),$$

where Γ is *Gamma function*.

The following result for powers holds:

THEOREM 1. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $p \geq 1$, $0 < \alpha < R$ and $x > 0$ with $\alpha x^p, \alpha x^{p-1} < R$, then*

$$0 \leq \frac{f(\alpha x^p)}{f(\alpha)} - \left[\frac{f(\alpha x)}{f(\alpha)} \right]^p \leq p \left[\frac{f(\alpha x^p)}{f(\alpha)} - \frac{f(\alpha x^{p-1})}{f(\alpha)} \frac{f(\alpha x)}{f(\alpha)} \right]. \tag{2.3}$$

Moreover, if $0 < x \leq 1$, then

$$\begin{aligned} 0 &\leq \frac{f(\alpha x^p)}{f(\alpha)} - \left[\frac{f(\alpha x)}{f(\alpha)} \right]^p \\ &\leq p \left[\frac{f(\alpha x^p)}{f(\alpha)} - \frac{f(\alpha x^{p-1})}{f(\alpha)} \frac{f(\alpha x)}{f(\alpha)} \right] \\ &\leq \frac{1}{2^p} \left(\frac{f(\alpha x^{2(p-1)})}{f(\alpha)} - \left[\frac{f(\alpha x^{p-1})}{f(\alpha)} \right]^2 \right)^{1/2} \leq \frac{1}{4^p} \end{aligned} \tag{2.4}$$

and

$$\begin{aligned} 0 &\leq \frac{f(\alpha x^p)}{f(\alpha)} - \left[\frac{f(\alpha x)}{f(\alpha)} \right]^p \\ &\leq p \left[\frac{f(\alpha x^p)}{f(\alpha)} - \frac{f(\alpha x^{p-1})}{f(\alpha)} \frac{f(\alpha x)}{f(\alpha)} \right] \\ &\leq \frac{1}{2^p} \left(\frac{f(\alpha x^2)}{f(\alpha)} - \left[\frac{f(\alpha x)}{f(\alpha)} \right]^2 \right)^{1/2} \leq \frac{1}{4^p}. \end{aligned} \tag{2.5}$$

Proof. If we write the inequality (1.1) for the convex function $\Phi : [0, \infty) \rightarrow [0, \infty)$, $\Phi(x) = x^p$, $p \geq 1$, then we have

$$\begin{aligned} 0 &\leq \sum_{i=1}^n w_i x_i^p - \left(\sum_{i=1}^n w_i x_i \right)^p \\ &\leq p \left(\sum_{i=1}^n w_i x_i^p - \sum_{i=1}^n w_i x_i^{p-1} \sum_{i=1}^n w_i x_i \right) \end{aligned} \quad (2.6)$$

for any $w_i, x_i \geq 0$ ($i = 1, \dots, n$) with $\sum_{i=1}^n w_i = 1$.

If $0 < \alpha < R$ and $k \geq 1$, then by (2.6) we have

$$\begin{aligned} 0 &\leq \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^p)^j - \left(\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j x^j \right)^p \\ &\leq p \left[\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^p)^j \right. \\ &\quad \left. - \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^{p-1})^j \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j x^j \right]. \end{aligned} \quad (2.7)$$

Since all series whose partial sums involved in the inequality (2.7) are convergent, then by letting $k \rightarrow \infty$ in (2.7) we deduce (2.3).

Now, if $x_i \in [m, M] \subset [0, \infty)$, ($i = 1, \dots, n$), then by (1.4) for the convex function $\Phi : [0, \infty) \rightarrow [0, \infty)$, $\Phi(x) = x^p$, $p \geq 1$ we have

$$\begin{aligned} 0 &\leq \sum_{i=1}^n w_i x_i^p - \left(\sum_{i=1}^n w_i x_i \right)^p \\ &\leq p \left(\sum_{i=1}^n w_i x_i^p - \sum_{i=1}^n w_i x_i^{p-1} \sum_{i=1}^n w_i x_i \right) \\ &\leq \frac{1}{2} p (M - m) \sum_{i=1}^n w_i \left| x_i^{p-1} - \sum_{j=1}^n w_j x_j^{p-1} \right| \\ &\leq \frac{1}{2} p (M - m) \left[\sum_{i=1}^n w_i x_i^{2(p-1)} - \left(\sum_{i=1}^n w_i x_i^{p-1} \right)^2 \right]^{1/2} \\ &\leq \frac{1}{4} p (M - m) (M^{p-1} - m^{p-1}). \end{aligned} \quad (2.8)$$

If $0 < x \leq 1$, then $0 < x^j \leq 1$ for $j = 0, \dots, k$, and by (2.8), we have

$$\begin{aligned}
 0 &\leq \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^p)^j - \left(\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j x^j \right)^p \\
 &\leq p \left[\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^p)^j \right. \\
 &\quad \left. - \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^{p-1})^j \frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j x^j \right] \\
 &\leq \frac{1}{2} p \\
 &\quad \times \left[\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j [x^{2(p-1)}]^j - \left(\frac{1}{\sum_{j=0}^k a_j \alpha^j} \sum_{j=0}^k a_j \alpha^j (x^{p-1})^j \right)^2 \right]^{1/2} \\
 &\leq \frac{1}{4} p. \tag{2.9}
 \end{aligned}$$

Since all series whose partial sums involved in the inequality (2.9) are convergent, then by letting $k \rightarrow \infty$ in (2.9) we deduce (2.4).

Now, if $x_i \in [m, M] \subset [0, \infty)$, ($i = 1, \dots, n$), then by (1.5) for the convex function $\Phi: [0, \infty) \rightarrow [0, \infty)$, $\Phi(x) = x^p$, $p \geq 1$ we have

$$\begin{aligned}
 0 &\leq \sum_{i=1}^n w_i x_i^p - \left(\sum_{i=1}^n w_i x_i \right)^p \\
 &\leq p \left(\sum_{i=1}^n w_i x_i^p - \sum_{i=1}^n w_i x_i^{p-1} \sum_{i=1}^n w_i x_i \right) \\
 &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \sum_{i=1}^n w_i \left| x_i - \sum_{j=1}^n w_j x_j \right| \\
 &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \left[\sum_{i=1}^n w_i x_i^2 - \left(\sum_{i=1}^n w_i x_i \right)^2 \right]^{1/2} \\
 &\leq \frac{1}{4} p (M - m) (M^{p-1} - m^{p-1}). \tag{2.10}
 \end{aligned}$$

Finally, by utilizing a similar argument as above, we obtain the inequality (2.5). The details are omitted. \square

Remark 1. We observe that the second inequality in (2.3) is equivalent to

$$\frac{f(\alpha x)}{f(\alpha)} \left(p \frac{f(\alpha x^{p-1})}{f(\alpha)} - \left[\frac{f(\alpha x)}{f(\alpha)} \right]^{p-1} \right) \leq (p-1) \frac{f(\alpha x^p)}{f(\alpha)} \quad (2.11)$$

or to

$$f(\alpha x) \left(p f(\alpha x^{p-1}) [f(\alpha)]^{p-2} - [f(\alpha x)]^{p-1} \right) \leq (p-1) f(\alpha x^p) [f(\alpha)]^{p-1}, \quad (2.12)$$

provided that $p \geq 1$, $0 < \alpha < R$ and $x > 0$ with $\alpha x^p, \alpha x^{p-1} < R$.

Moreover, if $0 < x \leq 1$, then from (2.4) we have

$$\left[\frac{f(\alpha x)}{f(\alpha)} \right]^p \leq \frac{f(\alpha x^p)}{f(\alpha)} \leq \frac{1}{4} p + \left[\frac{f(\alpha x)}{f(\alpha)} \right]^p. \quad (2.13)$$

Taking the power $1/p$ and using the inequality $(a+b)^{1/p} \leq a^{1/p} + b^{1/p}$, $p \geq 1$, we get

$$0 \leq [f(\alpha x^p)]^{1/p} [f(\alpha)]^{1-\frac{1}{p}} - f(\alpha x) \leq \frac{1}{4^{1/p}} p^{1/p} f(\alpha). \quad (2.14)$$

COROLLARY 1. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $u, v > 0$ with $v^p \leq u^q < R$, then

$$\left[\frac{f(uv)}{f(u^q)} \right]^p \leq \frac{f(v^p)}{f(u^q)} \leq \frac{1}{4} p + \left[\frac{f(uv)}{f(u^q)} \right]^p \quad (2.15)$$

and

$$0 \leq [f(v^p)]^{1/p} [f(u^q)]^{1/q} - f(uv) \leq \frac{1}{4^{1/p}} p^{1/p} f(u^q). \quad (2.16)$$

Proof. Follows by taking into (2.13) and (2.14) $\alpha = u^q$ and $x = \frac{v}{u^{q/p}}$. The details are omitted. \square

EXAMPLE 1.

a) If we write the inequalities (2.4) and (2.5) for the function $\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$, $z \in D(0, 1)$, then we have

$$\begin{aligned} 0 &\leq \frac{1-\alpha}{1-\alpha x^p} - \left(\frac{1-\alpha}{1-\alpha x} \right)^p \\ &\leq p \left[\frac{1-\alpha}{1-\alpha x^p} - \frac{(1-\alpha)^2}{(1-\alpha x^{p-1})(1-\alpha x)} \right] \\ &\leq \frac{1}{2} p \left[\frac{1-\alpha}{1-\alpha x^{2(p-1)}} - \left(\frac{1-\alpha}{1-\alpha x^{p-1}} \right)^2 \right]^{1/2} \leq \frac{1}{4} p \end{aligned} \quad (2.17)$$

and

$$\begin{aligned}
 0 &\leq \frac{1-\alpha}{1-\alpha x^p} - \left(\frac{1-\alpha}{1-\alpha x} \right)^p \\
 &\leq p \left[\frac{1-\alpha}{1-\alpha x^p} - \frac{(1-\alpha)^2}{(1-\alpha x^{p-1})(1-\alpha x)} \right] \\
 &\leq \frac{1}{2} p \left[\frac{1-\alpha}{1-\alpha x^2} - \left(\frac{1-\alpha}{1-\alpha x} \right)^2 \right]^{1/2} \leq \frac{1}{4} p
 \end{aligned} \tag{2.18}$$

for any $\alpha, x \in (0, 1)$ and $p \geq 1$.

b) If we write the inequalities (2.4) and (2.5) for the function $\exp z = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$, $z \in \mathbb{C}$, then we have

$$\begin{aligned}
 0 &\leq \exp[\alpha(x^p - 1)] - \exp[p\alpha(x - 1)] \\
 &\leq p \left[\exp[\alpha(x^p - 1)] - \exp[\alpha(x^{p-1} + x - 2)] \right] \\
 &\leq \frac{1}{2} p \left(\exp[\alpha(x^{2(p-1)} - 1)] - \exp[2\alpha(x^{p-1} - 1)] \right)^{1/2} \leq \frac{1}{4} p
 \end{aligned} \tag{2.19}$$

and

$$\begin{aligned}
 0 &\leq \exp[\alpha(x^p - 1)] - \exp[p\alpha(x - 1)] \\
 &\leq p \left[\exp[\alpha(x^p - 1)] - \exp[\alpha(x^{p-1} + x - 2)] \right] \\
 &\leq \frac{1}{2} p \left(\exp[\alpha(x^2 - 1)] - \exp[2\alpha(x - 1)] \right)^{1/2} \leq \frac{1}{4} p.
 \end{aligned} \tag{2.20}$$

for any $\alpha, p > 0$ and $x \in (0, 1)$.

3. Exponential inequalities

The following exponential inequality holds:

THEOREM 2. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $0 < \alpha < R$ and $x, \beta \in \mathbb{R}$ with $\alpha \exp(\beta x) < R$, then*

$$\begin{aligned}
 0 &\leq \frac{f(\alpha \exp(\beta x))}{f(\alpha)} - \exp \left[\frac{\alpha \beta x f'(\alpha)}{f(\alpha)} \right] \\
 &\leq \alpha \beta x \left[\frac{\exp(\beta x) f'(\alpha \exp(\beta x))}{f(\alpha)} - \frac{f(\alpha \exp(\beta x))}{f(\alpha)} \frac{f'(\alpha)}{f(\alpha)} \right].
 \end{aligned} \tag{3.1}$$

Moreover, if $x \leq 0$, $\beta > 0$ with $\exp(\beta x) < R$ and $0 < \alpha < R$, then

$$\begin{aligned}
 0 &\leq \frac{f(\alpha \exp(\beta x))}{f(\alpha)} - \exp\left[\frac{\alpha \beta x f'(\alpha)}{f(\alpha)}\right] \\
 &\leq \alpha \beta x \left[\frac{\exp(\beta x) f'(\alpha \exp(\beta x))}{f(\alpha)} - \frac{f(\alpha \exp(\beta x)) f'(\alpha)}{f(\alpha)^2} \right] \\
 &\leq \frac{1}{2} \beta |x| \left[\frac{\alpha [f'(\alpha) + \alpha f''(\alpha)]}{f(\alpha)} - \left(\frac{\alpha f'(\alpha)}{f(\alpha)} \right)^2 \right]^{1/2}. \tag{3.2}
 \end{aligned}$$

Proof. It is similar to the proof in Theorem 1 for the convex function $\Phi: \mathbb{R} \rightarrow [0, \infty)$, $\Phi(x) = \exp(\beta x)$ and the choice $x_j = jx$, $j \in \{0, \dots, k\}$. The details are omitted. \square

EXAMPLE 2.

- a) If we write the inequality (3.2) for the function $\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$, $z \in D(0, 1)$, then we have for $x \leq 0$, $\beta > 0$ and $0 < \alpha < 1$, that

$$\begin{aligned}
 0 &\leq \frac{1 - \alpha}{1 - \alpha \exp(\beta x)} - \exp\left(\frac{\alpha \beta x}{1 - \alpha}\right) \\
 &\leq \alpha \beta x \left[\frac{(1 - \alpha) \exp(\beta x)}{(1 - \alpha \exp(\beta x))^2} - \frac{1}{1 - \alpha \exp(\beta x)} \right] \leq \frac{1}{2} \frac{\beta |x| \alpha^{1/2}}{1 - \alpha}. \tag{3.3}
 \end{aligned}$$

- b) If we write the inequality (3.1) for the function $\exp z = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$, $z \in \mathbb{C}$, then we have

$$\begin{aligned}
 0 &\leq \exp\left(\alpha [\exp(\beta x) - 1]\right) - \exp(\alpha \beta x) \\
 &\leq \alpha \beta x \left[\exp\left(\alpha [\exp(\beta x) - 1] + \beta x\right) - \exp\left(\alpha [\exp(\beta x) - 1]\right) \right] \tag{3.4}
 \end{aligned}$$

for any $\alpha > 0$ and $x \leq 0$, $\beta > 0$.

4. Logarithmic inequalities

The following logarithmic inequality holds:

THEOREM 3. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $0 < \alpha < R$, $p > 0$ and $x > 0$ with $\alpha x^p, \alpha x^{-p} < R$, then

$$0 \leq \ln\left(\frac{f(\alpha x^p)}{f(\alpha)}\right) - p \frac{\alpha f'(\alpha)}{f(\alpha)} \ln x \leq \frac{f(\alpha x^p)}{f(\alpha)} \frac{f(\alpha x^{-p})}{f(\alpha)} - 1. \tag{4.1}$$

Moreover, if $0 < x \leq 1$ with $\alpha x^p, \alpha x^{-p} < R$, then

$$\begin{aligned} 0 &\leq \ln \left(\frac{f(\alpha x^p)}{f(\alpha)} \right) - p \frac{\alpha f'(\alpha)}{f(\alpha)} \ln x \leq \frac{f(\alpha x^p)}{f(\alpha)} \frac{f(\alpha x^{-p})}{f(\alpha)} - 1 \\ &\leq \frac{1}{2} \left[\frac{f(\alpha x^{-2p})}{f(\alpha)} - \left(\frac{f(\alpha x^{-p})}{f(\alpha)} \right)^2 \right]^{1/2}. \end{aligned} \quad (4.2)$$

Proof. It is similar to the proof in Theorem 1 for the convex function $\Phi: (0, \infty) \rightarrow \mathbb{R}$, $\Phi(x) = -\ln x$ and the choice $x_j = (x^p)^j$, $j \in \{0, \dots, k\}$. The details are omitted. \square

COROLLARY 2. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $u, v > 0$ with $v^p, u^q, \frac{u^{2q}}{v^p} < R$, then

$$0 \leq \ln \left(\frac{f(v^p)}{f(u^q)} \right) - \frac{u^q f'(u^q)}{f(u^q)} \ln \frac{v^p}{u^q} \leq \frac{f(v^p)}{f(u^q)} \frac{f\left(\frac{u^{2q}}{v^p}\right)}{f(u^q)} - 1. \quad (4.3)$$

If $v^p \leq u^q < R$ and $\frac{u^{2q}}{v^p}, \frac{u^{3q}}{v^{2p}}$, then

$$\begin{aligned} 0 &\leq \ln \left(\frac{f(v^p)}{f(u^q)} \right) - \frac{u^q f'(u^q)}{f(u^q)} \ln \frac{v^p}{u^q} \leq \frac{f(v^p)}{f(u^q)} \frac{f\left(\frac{u^{2q}}{v^p}\right)}{f(u^q)} - 1 \\ &\leq \frac{1}{2} \left[\frac{f\left(\frac{u^{3q}}{v^{2p}}\right)}{f(u^q)} - \left(\frac{f\left(\frac{u^{2q}}{v^p}\right)}{f(u^q)} \right)^2 \right]^{1/2}. \end{aligned} \quad (4.4)$$

Proof. Follows by taking $\alpha = u^q$ and $x = \frac{v}{u^{q/p}}$ into (4.1) and (4.2). The details are omitted. \square

EXAMPLE 3.

- a) If we write the inequality (4.2) for the function $\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$, $z \in D(0, 1)$, then we have for $0 < \alpha, x < 1$ and $p > 0$ that

$$\begin{aligned} 0 &\leq \ln \left(\frac{1-\alpha}{1-\alpha x^p} \right) - \frac{p\alpha}{1-\alpha} \ln x \\ &\leq \frac{(1-\alpha)^2}{(1-\alpha x^p)(1-\alpha x^{-p})} - 1 \\ &\leq \frac{1}{2} \left[\frac{1-\alpha}{1-\alpha x^{-2p}} - \left(\frac{1-\alpha}{1-\alpha x^{-p}} \right)^2 \right]^{1/2}. \end{aligned} \quad (4.5)$$

- b) If we write the inequality (4.1) for the function $\exp z = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$, $z \in \mathbb{C}$, then we have

$$0 \leq \alpha(x^p - 1) - p\alpha \ln x \leq \exp[\alpha(x^p + x^{-p} - 2)] - 1 \quad (4.6)$$

for $\alpha, p, x > 0$.

The following logarithmic inequality also holds:

THEOREM 4. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $0 < \alpha < R$, $p > 0$ and $x > 0$ with $\alpha x^p < R$, then*

$$\begin{aligned} 0 &\leq \frac{p\alpha x^p f'(\alpha x^p)}{f(\alpha)} \ln x - \frac{f(\alpha x^p)}{f(\alpha)} \ln \left(\frac{f(\alpha x^p)}{f(\alpha)} \right) \\ &\leq p\alpha \left[\frac{x^p f'(\alpha x^p)}{f(\alpha)} - \frac{f(\alpha x^p) f'(\alpha)}{f(\alpha)^2} \right] \ln x, \end{aligned} \quad (4.7)$$

or, equivalently

$$\frac{p\alpha f'(\alpha)}{f(\alpha)} \ln x \leq \ln \left(\frac{f(\alpha x^p)}{f(\alpha)} \right)$$

i.e., the first inequality in (4.1).

Moreover, if $0 < x \leq 1$ we also have

$$\begin{aligned} 0 &\leq \frac{p\alpha x^p f'(\alpha x^p)}{f(\alpha)} \ln x - \frac{f(\alpha x^p)}{f(\alpha)} \ln \left(\frac{f(\alpha x^p)}{f(\alpha)} \right) \\ &\leq p\alpha \left[\frac{x^p f'(\alpha x^p)}{f(\alpha)} - \frac{f(\alpha x^p) f'(\alpha)}{f(\alpha)^2} \right] \ln x \\ &\leq \frac{1}{2^p} |\ln x| \left[\frac{\alpha [f'(\alpha) + \alpha f''(\alpha)]}{f(\alpha)} - \left(\frac{\alpha f'(\alpha)}{f(\alpha)} \right)^2 \right]^{1/2}. \end{aligned} \quad (4.8)$$

Proof. It is similar to the proof in Theorem 1 for the convex function $\Phi: (0, \infty) \rightarrow \mathbb{R}$, $\Phi(x) = x \ln x$, and the choice $x_j = (x^p)^j$, $j \in \{0, \dots, k\}$. The details are omitted. \square

COROLLARY 3. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with nonnegative coefficients and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$.*

If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $u, v > 0$ with $v^p \leq u^q < R$, then

$$\begin{aligned} 0 &\leq \frac{f(v^p)}{f(u^q)} \ln \left(\frac{f(u^q)}{f(v^p)} \right) - \frac{v^p f'(v^p)}{f(u^q)} \ln \left(\frac{u^q}{v^p} \right) \\ &\leq u^q \left[\frac{f(v^p)}{f(u^q)} \frac{f'(u^q)}{f(u^q)} - \frac{\frac{v^p}{u^q} f'(v^p)}{f(u^q)} \right] \ln \left(\frac{u^q}{v^p} \right) \\ &\leq \frac{1}{2} \left[\frac{u^q [f'(u^q) + u^q f''(u^q)]}{f(u^q)} - \left(\frac{u^q f'(u^q)}{f(u^q)} \right)^2 \right]^{1/2} \ln \left(\frac{u^q}{v^p} \right). \end{aligned} \quad (4.9)$$

EXAMPLE 4. a) If we write the inequality (4.8) for the function $\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$, $z \in D(0, 1)$, then we have for $\alpha, x \in (0, 1)$ and $p > 0$ that

$$\begin{aligned} 0 &\leq \frac{p\alpha x^p (1-\alpha)}{(1-\alpha x^p)^2} \ln x - \frac{1-\alpha}{(1-\alpha x^p)} \ln \left(\frac{1-\alpha}{1-\alpha x^p} \right) \\ &\leq p\alpha \left[\frac{x^p (1-\alpha)}{(1-\alpha x^p)^2} - \frac{1}{1-\alpha x^p} \right] \ln x \\ &\leq \frac{1}{2} \frac{p\alpha^{1/2}}{1-\alpha} |\ln x|. \end{aligned} \quad (4.10)$$

b) If we write the inequality (4.8) for the function $\exp z = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$, $z \in \mathbb{C}$, then we have

$$\begin{aligned} 0 &\leq [p\alpha x^p \ln x - \alpha(x^p - 1)] \exp[\alpha(x^p - 1)] \\ &\leq p\alpha(x^p - 1) \exp[\alpha(x^p - 1)] \ln x \\ &\leq \frac{1}{2} p |\ln x| \alpha^{1/2} \end{aligned} \quad (4.11)$$

for $x \in (0, 1)$ and $\alpha, p > 0$.

Acknowledgements. The author would like to thank the anonymous referee for valuable recommendations that have been implemented in the final version of this paper.

REFERENCES

- [1] CERONE, P.—DRAGOMIR, S. S.: *A refinement of the Grüss inequality and applications*, Tamkang J. Math. **38** (2007), 37–49. Preprint *RGMA Res. Rep. Coll.* **5** (2002), Art. 14. <http://rgmia.org/papers/v5n2/RGIApp.pdf>.
- [2] CERONE, P.—DRAGOMIR, S. S.: *Some applications of de Bruijn’s inequality for power series*, Integral Transforms Spec. Funct. **18** (2007), 387–396.

- [3] DRAGOMIR, S. S.: *Discrete Inequalities of the Cauchy-Bunyakovsky-Schwarz Type*, Nova Sci. Publ. Inc., Hauppauge, N.Y., 2004.
- [4] DRAGOMIR, S. S.—IONESCU, N. M.: *Some converse of Jensen's inequality and applications*, Rev. Anal. Numér. Théor. Approx. **23** (1994), 71–78.
- [5] IBRAHIM, A.—DRAGOMIR, S. S.: *Power series inequalities via Buzano's result and applications*, Integral Transforms Spec. Funct. **22** (2011), 867–878.
- [6] IBRAHIM, A.—DRAGOMIR, S. S.: *Power series inequalities via a refinement of Schwarz inequality*, Integral Transforms Spec. Funct. **23** (2012), 769–78.
- [7] IBRAHIM, A.—DRAGOMIR, S. S.: *A survey on Cauchy–Bunyakovsky–Schwarz inequality for power series*, in: *Analytic Number Theory, Approximation Theory, and Special Functions* (G. V. Milovanović and M. Th. Rassias, eds.), Springer, New York, 2013, pp. 247–295.
- [8] IBRAHIM, A.—DRAGOMIR, S. S.—DARUS, M.: *Some inequalities for power series with applications*, Integral Transforms Spec. Funct. **24** (2013), 364–376.
- [9] IBRAHIM, A.—DRAGOMIR, S. S.—DARUS, M.: *Power series inequalities related to Young's inequality and applications*, Integral Transforms Spec. Funct. **24** (2013), 700–714.
- [10] IBRAHIM, A.—DRAGOMIR, S. S.—DARUS, M.: *Power series inequalities via Young's inequality with applications*, J. Inequal. Appl. **2013**, Article ID 314, 13 pp.

Received March 23, 2015

Mathematics
School of Engineering and Science
Victoria University
PO Box 14428
Melbourne City, MC 8001
AUSTRALIA
E-mail: sever.dragomir@vu.edu.au

School of Computational and
Applied Mathematics
University of the Witwatersrand
Private Bag 3
Johannesburg 2050
SOUTH AFRICA