

THE RIEMANN DERANGEMENT THEOREM AND DIVERGENT PERMUTATIONS

ROMAN WITUŁA

ABSTRACT. In this paper, a strengthening of the Riemann Derangement Theorem, by selecting the appropriate permutation of \mathbb{N} from two families \mathfrak{DC} and \mathfrak{DD} of permutations of \mathbb{N} , is presented. The mentioned families are characterized in a natural way; their algebraic properties were investigated by the author in his previous works.

1. Introduction

Subjects referring to the Riemann Derangement Theorem are still popular because the theorem offers a great inspiration. The recent papers [8], [9] (see also [6]) can be set the good examples of this statement. Let us recall its content.

THE RIEMANN DERANGEMENT THEOREM. *Let $\sum a_n$ be a conditionally convergent real series. Then, for every nonempty and closed interval $I \subset \mathbb{R}^*$ (two points compactification of \mathbb{R}), there exists a permutation p of \mathbb{N} such that the set of limit points of series $\sum a_{p(n)}$, denoted by $\sigma_{a_{p(n)}}$, is equal to I .*

In the current paper, we also will return to this classical result by selecting the appropriate permutation p of \mathbb{N} for this theorem from two subfamilies \mathfrak{DC} and \mathfrak{DD} of family \mathfrak{P} of all permutations of \mathbb{N} , as being discussed by author (see [1], [2], [7]) earlier.

However, we first need to introduce a few essential concepts. For short, we write $A < B$ for any two nonempty subsets A and B of \mathbb{N} when $a < b$ for each $a \in A$ and $b \in B$.

We call a sequence $\{A_n\}$ of nonempty subsets of \mathbb{N} to be increasing if

$$A_n < A_{n+1} \quad \text{for every } n \in \mathbb{N}.$$

We say that a nonempty subset A of \mathbb{N} is a union of k mutually separated intervals (abbreviated to **MSI**), if there exist k intervals I_1, \dots, I_k of \mathbb{N} which form a partition of A and $\text{dist}(I_i, I_j) \geq 2$ for any distinct indices $i, j \leq k$.

A permutation $p \in \mathfrak{P}$ is said to be divergent permutation if there is a conditionally convergent series $\sum a_n$ of real terms such that the p -rearranged series $\sum a_{p(n)}$ is divergent. The family of all divergent permutations will be denoted by \mathfrak{D} . Elements of family $\mathfrak{C} := \mathfrak{P} \setminus \mathfrak{D}$ are called the convergent permutations. We note that if $p \in \mathfrak{C}$ then, for every conditionally convergent series $\sum a_n$, the p -rearranged series $\sum a_{p(n)}$ is also convergent.

Family \mathfrak{D} will be naturally partitioned onto two subfamilies \mathfrak{DC} and \mathfrak{DD} , depending on the condition whether, for a given $p \in \mathfrak{D}$, we have $p^{-1} \in \mathfrak{C}$ or $p^{-1} \in \mathfrak{D}$, respectively. Some algebraic properties of those families were investigated by author [1]; among others, the following two relations

$$\mathfrak{DC} \circ \mathfrak{DC} = \mathfrak{DC} \quad \text{and} \quad \mathfrak{DD} \circ \mathfrak{DC} = \mathfrak{DC} \circ \mathfrak{DD} = \mathfrak{DC} \cup \mathfrak{DD}$$

were shown. The sign “ \circ ” denotes the composition of sets of permutations of \mathbb{N} here. In the current paper, we will add some other properties of \mathfrak{DC} and \mathfrak{DD} , of analytical nature, connected with the Riemann Derangement Theorem.

2. Main result

The two following dual combinatoric characterizations of convergent and divergent permutations are known [3], [4], [5] and will be explored in the proof of our main result.

THEOREM 2.1. *A permutation $p \in \mathfrak{P}$ is a convergent permutation if and only if there exists a positive integer k such that the set $p(I)$ is a union of at most k MSI for every interval I of \mathbb{N} . The minimal positive integer k with this property will be denoted by $\mathbf{c}(p)$.*

THEOREM 2.2. *A permutation $p \in \mathfrak{P}$ is a divergent permutation if and only if, for every positive integer n , there exists an interval I of \mathbb{N} such that $p(I)$ is a union of at least n MSI.*

Now, we are ready to formulate and to prove the main result—a strengthened version of the Riemann Derangement Theorem (announced by author in [7]).

THEOREM 2.3. *Let $\sum a_n$ be a conditionally convergent series and let $I \subset \mathbb{R}^*$ be a nondegenerated closed interval in \mathbb{R}^* . Then there exists a permutation $p \in \mathfrak{DD}$ such that $\sigma a_{p(n)} = I$.*

If we assume additionally that $\sum a_n \in I$ or that interval I is of the form $[\alpha, +\infty]$ or $[-\infty, \beta]$, for $\alpha, \beta \in \mathbb{R}^$, $\alpha < +\infty$ and $\beta > -\infty$, then there exists*

a permutation $q \in \mathfrak{DC}$ such that $\mathbf{c}(q^{-1}) \leq 5$ and $\sigma a_{q(n)} = I$. In the case $\sum a_n \in I$, the permutation $q \in \mathfrak{DC}$ can be selected in such a way that $\mathbf{c}(q^{-1}) \leq 3$.

Proof. Let $\alpha := \sum a_n$, $A^+ := \{i \in \mathbb{N} : a_i \geq 0\}$ and $A^- := \mathbb{N} \setminus A^+$. Furthermore, let us put $a_n^+ := \max\{0, a_n\}$ and $a_n^- := a_n - a_n^+$ for each $n \in \mathbb{N}$.

In the presented proof, we use the notations: (a, b) , $(a, b]$, $[a, b)$ and $[a, b]$ for the intervals of \mathbb{R} , as well as for the intervals of \mathbb{N} . Which notation is appropriate at a given moment, it will always clearly follow from the considerations.

Let $\beta, \gamma \in \mathbb{R}$, $\beta \leq \alpha \leq \gamma$. First, we define a permutation $q \in \mathfrak{DC}$ such that $\sigma a_{q(n)} := [\beta, \gamma]$ but the initial part of the proof is applicable to the case $\beta < \alpha < \gamma$ only. For this purpose, let us determine $m \in \mathbb{N}$ such that

$$|a_n| < \min \left\{ \frac{1}{2}(\gamma - \alpha), \frac{1}{2}(\alpha - \beta) \right\} \quad (1)$$

and

$$\frac{1}{2}(\beta - \alpha) < \alpha - \sum_{k=1}^n a_k < \frac{1}{2}(\gamma - \alpha), \quad (2)$$

for each $n \in \mathbb{N}$, $n > m$.

Next, by induction, we find the increasing sequence $\{I_n : n \in \mathbb{N}_0\}$ of intervals of \mathbb{N} forming the partition of \mathbb{N} and satisfying, for every $n \in \mathbb{N}$, the following conditions:

$$\sum_{j \in J_{2n-1}} a_j + \sum_{i \in I_{2n-1}} a_i^+ \geq \gamma, \quad \text{while} \quad \sum_{j \in J_{2n-1}} a_j + \sum_{\substack{i \in I_{2n-1} \\ i \neq k_{2n-1}}} a_i^+ \leq \gamma \quad (3)$$

and

$$\sum_{j \in J_{2n}} a_j + \sum_{i \in I_{2n}} a_i^- \leq \beta, \quad \text{while} \quad \sum_{j \in J_{2n}} a_j + \sum_{\substack{i \in I_{2n} \\ i \neq k_{2n}}} a_i^- \geq \beta, \quad (4)$$

where $I_0 := [1, m]$, $J_n := \bigcup_{i=0}^{n-1} I_i$ and

$$k_n := \begin{cases} \max\{k \in I_n : a_k > 0\} & \text{for } n \in (2\mathbb{N} - 1), \\ \max(I_n \cap A^-) & \text{for } n \in 2\mathbb{N}. \end{cases}$$

Additionally, we set

$$L_{2n-1} := I_{2n-1} \cap A^+, \quad L_{2n} := I_{2n} \cap A^- \quad \text{and} \quad l_n := \text{card } L_n,$$

for each $n \in \mathbb{N}$. We note that $\lim_{n \rightarrow \infty} l_n = \infty$. Moreover, if L_n is a union of λ_n **MSI** then, keeping in mind the conditional convergence of series $\sum a_n$ as well as conditions (3) and (4), we easily verify that also $\lim_{n \rightarrow \infty} \lambda_n = \infty$.

Permutation q is defined to be an increasing map of sets:

$$I_0 \cup \bigcup_{n \in \mathbb{N}} [\min I_n, l_n + \min I_n] \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} [l_n + \min I_n, \max I_n]$$

onto sets:

$$I_0 \cup \bigcup_{n \in \mathbb{N}} L_n \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} (I_n \setminus L_n),$$

respectively. One can easily verify that we then have

$$q([\min I_n, l_n + \min I_n]) = L_n, \quad \text{for every } n \in \mathbb{N},$$

and the set $q^{-1}(I)$ is a union of at most three **MSI** for every interval I , i.e., $q \in \mathfrak{DC}$ and $\mathbf{c}(q^{-1}) \leq 3$. In turn, from (3) and (4), it follows that $\sigma a_{q(n)} = [\beta, \gamma]$. However, it is not difficult to see that the above proof readily applies to the cases $\alpha = \beta$ or $\alpha = \gamma$ (for a nondegenerated interval $[\beta, \gamma]$).

Our next goal is to construct a permutation $q \in \mathfrak{DC}$ such that $\sum a_{q(n)} = +\infty$. Let $\{I_n\}$ be an increasing sequence of intervals of \mathbb{N} such that $\bigcup I_n = \mathbb{N}$ and

$$\sum_{i \in I_n} a_i^+ \geq n, \quad (5)$$

for each $n \in \mathbb{N}$. Moreover, let

$$\lambda_i := \text{card}(I_i \cap A^+) \quad \text{and} \quad \mu_i := \text{card}(I_i \cap A^-), \quad i \in \mathbb{N};$$

and

$$J_{n+1} := \left(\sum_{i=1}^{n+2} \lambda_i + \sum_{i=1}^n \mu_i, \sum_{i=1}^{n+2} \lambda_i + \sum_{i=1}^{n+1} \mu_i \right] \quad \text{for each } n \in \mathbb{N}.$$

We define the permutation q as an increasing map of sets:

$$[1, \lambda_1 + \lambda_2] \cup \bigcup_{n \in \mathbb{N}} \left(\sum_{i=1}^{n+1} \lambda_i + \sum_{i=1}^n \mu_i, \sum_{i=1}^{n+2} \lambda_i + \sum_{i=1}^n \mu_i \right]$$

and

$$(\lambda_1 + \lambda_2, \lambda_1 + \lambda_2 + \mu_1] \cup \bigcup_{n=2}^{\infty} J_n$$

onto sets A^+ and A^- , respectively.

One can verify that the permutation q possesses the following properties. First, we note that set $q^{-1}(I)$ is a union of at most five **MSI** for each interval $I \subset \mathbb{N}$, that is $q^{-1} \in \mathfrak{C}$ and $\mathbf{c}(q^{-1}) \leq 5$. Next, from condition (5), we obtain the following estimation

$$\sum_{i=1}^{\max J_n} a_{q(i)} = \sum_{i < \min J_n} a_i + \sum_{i \in J_n} a_i^+ \geq n + \sum_{i < \min J_n} a_i, \quad \text{correct for every } n \in \mathbb{N}.$$

Furthermore, it follows from the definition of q that the sequence

$$\left\{ \sum_{i=1}^j a_{q(i)} : \min J_n < j \leq \max J_n \right\} \text{ is non-increasing for each } n \in \mathbb{N},$$

whereas the sequence

$$\left\{ \sum_{i=1}^j a_{q(i)} : \max J_n < j < \min J_{n+1} \right\} \text{ is increasing for each } n \in \mathbb{N}.$$

From the last three properties, we receive $\sum a_{q(n)} = +\infty$. In consequence, we have $q \in \mathfrak{DC}$.

Let $\beta \in \mathbb{R}$. We want to define a permutation $q \in \mathfrak{DC}$ such that $\sigma a_{q(n)} = [\beta, \infty]$. For this purpose, it is enough to choose by induction the increasing sequence $\{I_n\}$ of intervals of \mathbb{N} forming the partition of \mathbb{N} such that for each $n \in \mathbb{N}$ the following conditions are satisfied

$$\sum_{j \in J_{2n-1}} a_{q(j)} \geq \beta + n \quad \text{and} \quad \sum_{j \in J_{2n}} a_{q(j)} \leq \beta, \quad (6)$$

while

$$\sum_{\substack{j \in J_{2n} \\ j < \max J_{2n}}} a_{q(j)} \geq \beta,$$

where $J_n := \bigcup_{i=0}^n I_i$, whereas q is defined to be the increasing map of sets $\bigcup_{n \in \mathbb{N}} I_{2n-1}$ and $\bigcup_{n \in \mathbb{N}} I_{2n}$ onto sets A^+ and A^- , respectively. Then, it is easy to verify that $q^{-1} \in \mathfrak{C}$ and $\mathbf{c}(q^{-1}) \leq 5$, and, since from (6) it results that $\sigma a_{q(n)} = [\beta, \infty]$, we have $q \in \mathfrak{DC}$.

Definition of permutation $q \in \mathfrak{DC}$, such that $\sigma a_{q(n)} = \mathbb{R}^*$, is the same as in the previously discussed case of $\sigma a_{q(n)} = [\beta, \infty]$, $\beta \in \mathbb{R}$, only condition (6) needs to be changed. More precisely, condition (6) must be replaced by the following conditions:

$$\sum_{j \in J_{2n-1}} a_{q(j)} \geq n \quad \text{and} \quad \sum_{j \in J_{2n}} a_{q(j)} \leq -n,$$

for each $n \in \mathbb{N}$.

Definition of permutation $p \in \mathfrak{DD}$, which we will also make dependent on the form of set $\sigma a_{p(n)}$, must be preceded by some essential preparations. The main charge of those preparations lies in the appropriate selection of increasing sequence $\{I_n : n \in \mathbb{N}_0\}$ of the intervals of \mathbb{N} forming the partition of \mathbb{N} . We denote as s and t the increasing bijections of \mathbb{N} onto sets

$$\mathbf{D} := \bigcup_{n \in \mathbb{N}} I_{2n-1} \quad \text{and} \quad \mathbf{E} := \bigcup_{n \in \mathbb{N}_0} I_{2n},$$

respectively. We request the intervals I_{2n} , $n \in \mathbb{N}_0$, to be composed of the even number of elements and so that $\text{card } I_{2n} \rightarrow \infty$ for $n \rightarrow \infty$. Moreover, we demand

the series $\sum a_{s(n)} = \sum_{n \in \mathbf{D}} a_n$ and $\sum a_{t(n)} = \sum_{n \in \mathbf{E}} a_n$ both to be convergent, in addition, the first one should be conditionally convergent and the second series $\sum a_{t(n)}$ should be absolutely convergent (and therefore unsusceptible to permutations). Let $d := \sum a_{t(n)}$.

Restriction of permutation p to set \mathbf{E} will be defined in the same way in all of the cases discussed below. So, we take that

$$p(2i - 2 + \min I_{2n}) = i - 1 + \min I_{2n}$$

and

$$p(2i - 1 + \min I_{2n}) = i - 1 + \frac{1}{2} \text{card } I_{2n} + \min I_{2n},$$

for $i = 1, 2, \dots, \frac{1}{2} \text{card } I_{2n}$ and for each $n \in \mathbb{N}_0$. It implies that for each case we have $p^{-1} \in \mathfrak{D}$.

First, let us consider the case in which $\sigma a_{p(n)}$ is a nondegenerated closed interval. Let $a, b \in \mathbb{R}$, $a < b$ and $d < b$ (the last condition does not violate the generality of considerations because one can assume d to be any real number). Additionally, we assume that

$$|a_j| < \frac{1}{2}|b - a| \quad \text{for every } j \in \mathbf{D}. \quad (7)$$

We define by induction an increasing sequence $\{J_n\}$ of intervals of \mathbb{N} forming the partition of \mathbb{N} such that

$$\mu \left(\bigcup_{n \in \mathbb{N}} J_{2n-1} \right) = \mathbf{D} \cap A^+, \quad \mu \left(\bigcup_{n \in \mathbb{N}} J_{2n} \right) = \mathbf{D} \cap A^-,$$

$$\left\{ \begin{array}{l} \sum_{j \in K_{2n-1}} a_{\mu(j)} \geq b - d, \quad \text{while} \quad \sum_{\substack{j \in K_{2n-1} \\ j < \max K_{2n-1}}} a_{\mu(j)} \leq b - d, \\ \text{and} \\ \sum_{j \in K_{2n}} a_{\mu(j)} \leq a - d, \quad \text{while} \quad \sum_{\substack{j \in K_{2n} \\ j < \max K_{2n}}} a_{\mu(j)} \geq a - d, \end{array} \right. \quad (8)$$

for each $n \in \mathbb{N}$, where $K_n := \bigcup_{j=1}^n J_j$ and $\mu := ps$. The restriction of p to set \mathbf{D} is defined to be the increasing map of sets

$$s \left(\bigcup_{n \in \mathbb{N}} J_{2n-1} \right) \quad \text{and} \quad s \left(\bigcup_{n \in \mathbb{N}} J_{2n} \right)$$

onto sets $\mathbf{D} \cap A^+$ and $\mathbf{D} \cap A^-$, respectively. Then, what can be easily verified, we have $\sigma a_{p(n)} = [a, b]$ which implies, in particular, that $p \in \mathfrak{D}$.

THE RIEMANN DERANGEMENT THEOREM AND DIVERGENT PERMUTATIONS

Let $a \in \mathbb{R}$. For $\sigma a_{p(n)} = [a, \infty]$ or $\sigma a_{p(n)} = \{+\infty\}$, the construction of permutation p is almost analogous as shown before, only conditions (7) and (8) must be modified. So, for $\sigma a_{p(n)} = [a, \infty]$, $a \in \mathbb{R}$, instead of condition (7), we assume that $|a_j| < \frac{1}{2}$, for every $j \in \mathbf{D}$, whereas (8) is replaced, for each $n \in \mathbb{N}$, with the following condition

$$\left\{ \begin{array}{l} \sum_{j \in K_{2n-1}} a_{\mu(j)} \geq \max\{n, 2a\}, \quad \text{while} \quad \sum_{\substack{j \in K_{2n-1} \\ j < \max K_{2n-1}}} a_{\mu(j)} \leq \max\{n, 2a\}, \\ \text{and} \\ \sum_{j \in K_{2n}} a_{\mu(j)} \leq a, \quad \text{while} \quad \sum_{\substack{j \in K_{2n} \\ j < \max K_{2n}}} a_{\mu(j)} \geq a. \end{array} \right. \quad (9)$$

For the case $\sigma a_{p(n)} = \{+\infty\}$, we request that $|a_j| < 1$, for each $j \in \mathbf{D}$, whereas (8) is replaced, for each $n \in \mathbb{N}$, with the condition given below

$$\left\{ \begin{array}{l} \sum_{j \in K_{2n-1}} a_{\mu(j)} \geq 2n, \quad \text{while} \quad \sum_{\substack{j \in K_{2n-1} \\ j < \max K_{2n-1}}} a_{\mu(j)} \leq 2n, \\ \text{and} \\ \sum_{j \in K_{2n}} a_{\mu(j)} \leq n, \quad \text{while} \quad \sum_{\substack{j \in K_{2n} \\ j < \max K_{2n}}} a_{\mu(j)} \geq n. \end{array} \right. \quad (10)$$

□

Acknowledgements. The author is grateful to the reviewer’s valuable comments that improved the manuscript.

REFERENCES

- [1] WITUŁA, R.: *The algebraic properties of the convergent and divergent permutations*, Filomat (submitted).
- [2] WITUŁA, R.: *The family \mathfrak{F} of permutations of \mathbb{N}* , Positivity (submitted).
- [3] WITUŁA, R.—SŁOTA, D.—SEWERYN, R.: *On Erdős’ theorem for monotonic subsequences*, Demonstr. Math. **40** (2007), 239–259.
- [4] WITUŁA, R.: *On the set of limit points of the partial sums of series rearranged by a given divergent permutation*, J. Math. Anal. Appl. **362** (2010), 542–552.
- [5] LEVI, F. W.: *Rearrangement of convergent series*, Duke Math. J. **13** (1946), 579–585.
- [6] WITUŁA, R.—HETMANIOK, E.—SŁOTA, D.: *Some variation on the Riemann Derangement Theorem*, Tatra Mt. Math. Publ. (submitted).
- [7] WITUŁA, R.: *The Riemann theorem and divergent permutations*, Colloq. Math. **69** (1995), 275–287.
- [8] FILIPÓW, R.—SZUCA, P.: *Rearrangement of conditionally convergent series on a small set*, J. Math. Anal. Appl. **362** (2010), 64–71.

ROMAN WITUŁA

- [9] PRUS-WIŚNIEWSKI, F.: *Two refinements of the Riemann Derangement Theorem*.
In: Real Functions, Density Topology and Related Topics (M. Filipczak and E. Wagner-
-Bojakowska, eds.), Łódź University Press, Łódź, 2011, pp. 165–172.

Received July 4, 2011

*Institute of Mathematics
Silesian University of Technology
Kaszubska 23
PL-Gliwice 44-100
POLAND
E-mail: roman.witula@polsl.pl*