

## APPLICATION OF SURFACE AND VOLUME GEOMETRY ANALYSIS IN GEOSCIENCES

JAROSLAV HOFIERKA<sup>1</sup> and MARIÁN ZLOCHA<sup>2</sup>

<sup>1</sup>Department of Cartography, Geoinformatic and Remote Sensing

<sup>2</sup>Department of Economic Geology, Faculty of Sciences, Comenius University, Mlynská dolina, 842 15 Bratislava, Slovak Republic

There are many tasks in geosciences where geometry analysis is required to better understand a given phenomenon or process. Surface geometry analysis was originally developed in geomorphology where geometrical properties of relief significantly influence geomorphological processes (e.g. water erosion, landslides, etc.). From the geometrical point of view, this analysis is 2-dimensional because 2 independent variables (x, y) are used. The phenomenon is modelled using a bivariate function  $z = f(x, y)$ . However in geological applications, 3-dimensional cases are more obvious. Therefore, a trivariate function  $w = f(x, y, z)$  is used. The most important surface geometry analysis parameters expressed using partial derivatives of bivariate function  $z = f(x, y)$  are (Krcho 1990), (Mitášová & Hofierka 1992): slope, aspect, profile curvature, tangential curvature.

Derived parameters are, for example, flowlines, their length and density interpreted often as an upslope contributing area and often used in water erosion modelling. These parameters are important in landscape processes where relief is the main factor in the spatial differentiation of these processes. There are many examples of using them in geomorphology, landscape ecology, pedology, etc. (Moore et al. 1991). In volume geometry analysis, parameters analogical to 2-D parameters have been derived in (Hofierka et al. 1993):

-magnitude of gradient  $|gradw| = \sqrt{f_x^2 + f_y^2}$

$$\delta |gradw| / \delta n = \frac{f_x^2 f_{xx} + 2f_x f_y f_{xy} + 2f_y f_x f_{xy} + f_y^2 f_{yy} + f_x^2 f_{zz}}{f_x^2 + f_y^2 + f_z^2}$$

- Gaussian curvature  $G = \begin{vmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{vmatrix} \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix}^{-1}$

- mean curvature  $H = \frac{1}{3}(A + B + C) \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix}^{-1}$

$$A = \begin{vmatrix} h_{11} & h_{12} & h_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix} \quad B = \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ h_{21} & h_{22} & h_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix} \quad C = \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ h_{31} & h_{32} & h_{33} \end{vmatrix}$$

where  $g_{11} = 1 + f_x^2$      $g_{22} = 1 + f_y^2$      $g_{33} = 1 + f_z^2$   
 $g_{21} = g_{12} = f_x f_y$      $g_{31} = g_{13} = f_x f_z$      $g_{32} = g_{23} = f_y f_z$

$$h_{11} = \frac{f_{xx}}{\sqrt{1+f_x^2+f_y^2+f_z^2}} \quad h_{22} = \frac{f_{yy}}{\sqrt{1+f_x^2+f_y^2+f_z^2}} \quad h_{33} = \frac{f_{zz}}{\sqrt{1+f_x^2+f_y^2+f_z^2}}$$

$$h_{21} = h_{12} = \frac{f_{xy}}{\sqrt{1+f_x^2+f_y^2+f_z^2}} \quad h_{31} = h_{13} = \frac{f_{xz}}{\sqrt{1+f_x^2+f_y^2+f_z^2}}$$

$$h_{23} = h_{32} = \frac{f_{yz}}{\sqrt{1+f_x^2+f_y^2+f_z^2}}$$

3-D flowlines, their length and density.

Magnitude of gradient, curvatures,  $\delta |gradw| / \delta n$ , can be used as topological parameters characterizing the shape of investigated geological solids. Their interpretation and usefulness are in the 3-D case more complicated because of the lack of driving factors like relief. These parameters are more described in (Zlocha et al. 1993). 3-D flowlines, their length and density have clear interpretation especially in hydrogeology. For example, places of high densities should indicate zones of matter accumulation. The best way of computing 2-D and 3-D morphometrical parameters is using an appropriate interpolation function (e.g. regularized spline with tension derived in (Mitášová & Mitáš 1992)). Large datasets (tens of thousands points) can be computed in a real time only using high performance computers.

### References

- Hofierka J., Mitášová H. & Zlocha M., 1993: Volume geometry analysis and computation of 3-D flowlines: I. Theory. Manuscript.
- Krcho J., 1990: Morphometric analysis and the digital models of georelief. *VEDA*, Bratislava, 432 (in Slovak).
- Mitášová H. & Hofierka J., 1992: Interpolation by regularized spline with tension: II. Application to terrain modeling and surface geometry analysis. *Mathematical Geology* (in press).
- Mitášová H. & Mitáš L., 1992: Interpolation by Regularized Spline with Tension: I. Theory and Implementation. *Mathematical Geology* (in press).
- Moore I. D., Grayson R. B. & Ladson A. R., 1991: Digital terrain modelling: a review of hydrological, geomorphological and biological applications. *Hydrological Processes* 5, 3 - 30.
- Zlocha M., Hofierka J. & Mitášová H., 1993: Volume geometry analysis and computation of 3-D flowlines: II. Application. Manuscript.