

ON ASYMPTOTIC BEHAVIOUR OF SOLUTIONS OF FUNCTIONAL DIFFERENTIAL EQUATIONS

ROMAN KOPLATADZE

ABSTRACT. Sufficient (necessary and sufficient) conditions are given for a functional differential equation to have properties *A* and *B*.

Consider the equation

$$u^{(n)}(t) + F(u)(t) = 0, \quad (1)$$

where $F: C^{n-1}(\mathbb{R}_+; \mathbb{R}) \rightarrow L_{\text{loc}}(\mathbb{R}_+; \mathbb{R})$ is a continuous operator. Everywhere below we shall assume that a nondecreasing function $\sigma: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ exists such that $\lim_{t \rightarrow +\infty} \sigma(t) = +\infty$ and for any $t \in \mathbb{R}_+$

$$F(x)(t) = F(y)(t) \quad \text{if } x, y \in C^{n-1}(\mathbb{R}_+; \mathbb{R})$$

and

$$x(s) = y(s) \quad \text{for } s \geq \sigma(t).$$

For any $t_0 \in \mathbb{R}_+$ let M_{t_0} denote the set of $u \in C^{n-1}(\mathbb{R}_+; \mathbb{R})$ satisfying $u(t) \neq 0$ for $t \geq t^*$, where $t^* = \min\{t_0, \sigma(t_0)\}$. The following assumption will always be fulfilled: either

$$F(u)(t) u(t) \geq 0 \quad \text{for } t \geq t_0, \quad \text{for any } t_0 \in \mathbb{R}_+ \text{ and } u \in M_{t_0}, \quad (2)$$

or

$$F(u)(t) u(t) \leq 0 \quad \text{for } t \geq t_0, \quad \text{for any } t_0 \in \mathbb{R}_+ \text{ and } u \in M_{t_0}. \quad (3)$$

AMS Subject Classification (1991): 34K15.

Key words: functional differential equations, properties *A* and *B*, oscillation of solutions.

DEFINITION 1. Let $t_0 \in \mathbb{R}_+$. A function $u: [t_0, +\infty[\rightarrow \mathbb{R}$ is said to be the *proper solution* of the equation (1) if it is locally absolutely continuous up to the order $n - 1$ inclusively, there exists a function $\bar{u} \in C^{n-1}(\mathbb{R}_+; \mathbb{R})$ such that $\bar{u}(t) \equiv u(t)$ for $t \geq t_0$, almost everywhere on $[t_0, +\infty[$

$$\bar{u}^{(n)}(t) + F(\bar{u})(t) = 0$$

and

$$\sup\{|u(s)| : s \in [t, +\infty[\} > 0 \quad \text{for any } t \in [t_0, +\infty[.$$

DEFINITION 2. We say that the equation (1) *has the property A* provided any of its proper solutions is oscillatory if n is even and either is oscillatory or satisfies

$$|u^{(i)}(t)| \downarrow 0 \quad \text{as } t \uparrow +\infty \quad (i = 0, \dots, n-1) \quad (4)$$

if n is odd.

DEFINITION 3. We say that the equation (1) *has the property B* provided any of its proper solutions either is oscillatory or satisfies (4) or

$$|u^{(i)}(t)| \uparrow 0 \quad \text{as } t \uparrow +\infty \quad (i = 0, \dots, n-1) \quad (5)$$

if n is even and either is oscillatory or satisfies (5) if n is odd.

Conditions for an ordinary differential equation to have the properties A and B are studied well enough (see [1, 2] and references therein). The analogous problems for the equations with deviating arguments are investigated in [3, 4].

THEOREM 1. Let (2) ((3)) hold and let for any $t_0 \in \mathbb{R}_+$

$$|F(u)(t)| \geq \sum_{i=1}^m \int_{\sigma_i(t)}^{\bar{\sigma}_i(t)} |u(s)| d_s r_i(t, s) \quad \text{for } t \geq t_0, \quad u \in M_{t_0}, \quad (6)$$

where the measurable functions $r_i(t, s)$ ($i = 1, \dots, m$) are nondecreasing in s , $\sigma_i, \bar{\sigma}_i \in C(\mathbb{R}_+; \mathbb{R}_+)$, $\sigma_i(t) \leq \bar{\sigma}_i(t)$ ($i = 1, \dots, m$) for $t \geq 0$ and

$$\lim_{t \rightarrow +\infty} \frac{\sigma_i(t)}{t} > 0 \quad (i = 1, \dots, m). \quad (7)$$

Suppose, moreover, that there exists $\varepsilon > 0$ such that for any $l \in \{1, \dots, n-1\}$ and $\lambda \in [l-1, l[$ where $l+n$ is odd (even) the inequality

$$\lim_{t \rightarrow +\infty} t^{l-\lambda} \int_t^{+\infty} \xi^{n-l-1} \sum_{i=1}^m \int_{\sigma_i(\xi)}^{\bar{\sigma}_i(\xi)} s^\lambda d_s r_i(\xi, s) d\xi \geq \prod_{i=0, i \neq l}^{n-1} |\lambda - i| + \varepsilon$$

holds. Then the equation (1) has the property A (B).

THEOREM 2. Let (2), (6), (7) ((3), (6), (7)) hold. Suppose, moreover, that there exists $\varepsilon > 0$ such that for any $\lambda \in \bigcup_{k=0}^{(n-2)/2} [2k, 2k+1[$ ($\lambda \in \bigcup_{k=1}^{(n-2)/2} [2k-1, 2k[$) if n is even and for any $\lambda \in \bigcup_{k=1}^{(n-1)/2} [2k-1, 2k[$ ($\lambda \in \bigcup_{k=0}^{(n-3)/2} [2k, 2k+1[$) if n is odd the inequality

$$\lim_{t \rightarrow +\infty} t \int_t^{+\infty} \xi^{n-\lambda-2} \sum_{i=1}^m \int_{\sigma_i(\xi)}^{\bar{\sigma}_i(\xi)} s^\lambda d_s r_i(\xi, s) d\xi \geq \prod_{i=0}^{n-1} |\lambda - i| + \varepsilon$$

holds. Then the equation (1) has the property A (B).

COROLLARY 1. Let (2), (7) ((3), (7)) hold and let for any $t_0 \in \mathbb{R}_+$

$$|F(u)(t)| \geq \sum_{i=1}^m p_i(t) |u(\sigma_i(t))| \quad \text{for } t \geq t_0, \quad u \in M_{t_0},$$

where $p_i \in L_{\text{loc}}(\mathbb{R}_+; \mathbb{R}_+)$, $\sigma_i \in C(\mathbb{R}_+; \mathbb{R}_+)$, $\sigma_i(t) \leq t$ ($i = 1, \dots, m$). Suppose, moreover, that there exists $\varepsilon > 0$ such that for any $\lambda \in [n-2, n-1[$ (for any $\lambda \in [n-3, n-2[\cup [0, 1[$ if n is odd and for any $\lambda \in [n-3, n-2[$ of n is even) the inequality

$$\lim_{t \rightarrow +\infty} t \int_t^{+\infty} \xi^{n-\lambda-2} \sum_{i=1}^m p_i(\xi) \sigma_i \sigma_i^\lambda(\xi) d\xi \geq \prod_{i=0}^{n-1} |\lambda - i| + \varepsilon$$

holds. Then the equation (1) has the property A (B).

COROLLARY 2. Let (2) ((3)) hold and let for any $t_0 \in \mathbb{R}_+$,

$$|F(u)(t)| \geq \frac{c}{t^{n+1}} \int_{\alpha t}^{\bar{\alpha} t} |u(s)| ds \quad \text{for } t \geq t_0, \quad u \in M_{t_0},$$

where $0 < \alpha < \bar{\alpha}$, and

$$c > \max \left\{ -(\lambda+1)\lambda(\lambda-1) \cdots (\lambda-n+1)(\bar{\alpha}^{\lambda+1} - \alpha^{\lambda+1}) : \lambda \in [0, n-1] \right\} \\ (c > \max \left\{ (\lambda+1)\lambda(\lambda-1) \cdots (\lambda-n+1)(\bar{\alpha}^{\lambda+1} - \alpha^{\lambda+1}) : \lambda \in [0, n-1] \right\}) \quad (8)$$

Then the equation (1) has the property A (B).

COROLLARY 3. Let $\alpha > 0$ and $c \in]0, +\infty[$ ($c \in]-\infty, 0[$). Then the condition

$$c > \max\{-\alpha^{-\lambda}\lambda(\lambda-1)\cdots(\lambda-n+1): \lambda \in [0, n-1]\} \\ (c < -\max\{\alpha^{-\lambda}\lambda(\lambda-1)\cdots(\lambda-n+1): \lambda \in [0, n-1]\})$$

is necessary and sufficient for the equation

$$u^{(n)}(t) + \frac{c}{t^n} u(\alpha t) = 0$$

to have the property $A(B)$.

COROLLARY 4. Let $0 < \alpha < \bar{\alpha}$ and $c \in]0, +\infty[$ ($c \in]-\infty, 0[$). Then the condition (8)

$$(c < -\max\{(\lambda+1)\lambda(\lambda-1)\cdots(\lambda-n+1)(\bar{\alpha}^{\lambda+1} - \alpha^{\lambda+1})^{-1}: \lambda \in [0, n-1]\})$$

is necessary and sufficient for the equation

$$u^{(n)}(t) + \frac{c}{t^{n+1}} \int_{\alpha t}^{\bar{\alpha} t} u(s) ds = 0$$

to have the property $A(B)$.

REFERENCES

- [1] KIGURADZE, I. T.—ČANTURIJA, T. A.: *Asymptotic Properties of Solutions of Nonautonomous Ordinary Differential Equations*, Moscow, 1990. (Russian)
- [2] KONDRÁTEV, V. A.: *On oscillation of solutions of the equation $y^{(n)} + P(x)y = 0$* , Trudy Moskov. Mat. Obshch. **10** (1961), 419–436. (Russian)
- [3] KOPLATADZE, R. G.—ČANTURIJA, T. A.: *On Oscillatory Properties of Differential Equations with Deviating Argument*, Tbilisi, 1977. (Russian)
- [4] KOPLATADZE, R. G.: *On differential equations with deviating argument having properties A and B*, Differentsialnye Uravneniya **V25** N11 (1989), 1897–1909. (Russian)

Received October 7, 1993

*I. N. Vekua Institute of Applied Mathematics
University str. 2
380043 Tbilisi
REPUBLIC OF GEORGIA
E-mail: kig@imath.kheta.georgia.su*